

On the criteria of $\mathcal{D}$ -planarity of a tree.

Polulyakh E., Yurchuk I.

Institute of Mathematics of Ukrainian national academy of sciences, Kyiv

Abstract. Let T be a tree with a fixed subset of vertices V^* such that there is a cyclic order C on it and all terminal vertices are contained in this set. Let $D^2 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$ be a closed oriented 2-dimensional disk. The tree T is called $\mathcal{D}$ -planar if there exists an embedding $\varphi : T \rightarrow \mathbb{R}^2$ which satisfies the following conditions $\varphi(T) \subseteq D^2$, $\varphi(T) \cap \partial D^2 = \varphi(V^*)$ and if $\#V^* \geq 3$ then a cyclic order $\varphi(C)$ of $\varphi(V^*)$ coincides with a cyclic order which is generated by the orientation of $\partial D^2 \cong S^1$.

We obtain a necessary and sufficient condition for T to be $\mathcal{D}$ -planar.

Keywords. $\mathcal{D}$ -planar tree, a cyclic order, a convenient relation.

INTRODUCTION.

It is known that any tree is planar (i.e., can be embedded into a plane), more over, in [7] it was proved that any rooted tree with n vertices can be embedded as a plane spanning tree on n points of a plane, with the root being mapped onto an arbitrary specified point of them. There are similar results for rooted star forests [8]. Many authors were interested in different types of embedding of trees into a disk with extra conditions. For example, in [9, 10] author used a special type of embedding of tree into a disk for constructing a Morse function on it.

In this paper we will consider an embedding of a tree T into D^2 (i.e., a closed oriented 2-dimensional disk) such that a fixed subset of its vertices V^* which contains all terminal vertices maps to a boundary of D^2 in a special way and $T \setminus V^*$ maps into $IntD^2$.

The existence of such embedding is a part of a solution of one topological problem: the realization of a finite graph as invariant of pseudoharmonic functions defined on a disk. Namely, a continuous function $f : D^2 \rightarrow \mathbb{R}$ is called a pseudoharmonic function if $f|_{\partial D^2}$ is a continuous function with a finite number of local extrema and $f|_{IntD^2}$ has a finite number of critical points and each of them is a saddle point (in the neighborhood of it f has a representation as $Rez^n + const$, $z = x + iy$ and $n \geq 2$). In [6] the topological invariant of such functions is constructed, in particular, it consists of a cycle γ with special properties and the complement to it is a disjoint union of trees. The criteria of topological equivalence of such functions is formulated in terms of their invariant.

In Section 1 we will describe some properties of a $\mathcal{D}$ -planar tree. They will be useful for the proof of Theorem 2.1 which gives the criteria of $\mathcal{D}$ -planarity of a tree. In this section we also study some properties of different types of relations on finite sets.

In Section 2 the criteria of $\mathcal{D}$ -planarity of a tree will be proved. Its proof has a topological nature.

We are sincerely grateful to V.V. Sharko, S.I. Maksymenko and I. Yu. Vlasenko for useful discussions.

1. THE UTILITY RESULTS.

1.1. Properties of trees embedded into two-dimensional disk. Let T be a tree with a set of vertices V and a set of edges E . Suppose that T is non degenerated (has at least one edge). Denote by V_{ter} a set of all vertices of T such that their degree equals to 1. Let us assume that for a subset $V^* \subseteq V$ the following condition holds true

$$(1) \quad V_{ter} \subseteq V^* .$$

Let also $\varphi : T \rightarrow \mathbb{R}^2$ is an embedding such that

$$(2) \quad \varphi(T) \subseteq D^2, \quad \varphi(T) \cap \partial D^2 = \varphi(V^*) .$$

Lemma 1.1. *A set $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$ has a finite number of connected components*

$$U_0 = \mathbb{R}^2 \setminus D^2, U_1, \dots, U_m ,$$

and for every $i \in \{1, \dots, m\}$ a set U_i is an open disk and is bounded by a simple closed curve

$$\partial U_i = L_i \cup \varphi(P(v_i, v'_i)), \quad L_i \cap \varphi(P(v_i, v'_i)) = \{\varphi(v_i), \varphi(v'_i)\}$$

where L_i is an arc of ∂D^2 such that the vertices $\varphi(v_i)$ and $\varphi(v'_i)$ are its endpoints, and $\varphi(P(v_i, v'_i))$ is an image of the unique path $P(v_i, v'_i)$ in T which connects v_i and v'_i .

Proof. We prove lemma by an induction on the number of elements of the set V^* . Denote by $\sharp A$ a number of elements of a set A .

Let us remark that $\sharp V^* \geq 2$ since $V_{ter} \subseteq V^*$ and $\sharp V_{ter} \geq 2$ for a non degenerated tree T (it is easily verified by induction on the number of vertices).

Base of induction. Let $\sharp V^* = 2$. From what was said above it follows that $V^* = V_{ter}$. So, a tree satisfies a condition $\sharp V_{ter} = 2$. For such trees it is easy to prove by induction on the number of vertices that every vertex of $V \setminus V_{ter}$ has degree 2. In other words it is adjacent to two edges.

If a tree is considered as CW-complex (i.e. 0-cells are its vertices and 1-cells are its edges), then a topological space T is homeomorphic to a segment with a set of the endpoints which coincides with $V^* = V_{ter}$.

Let $\varphi(T)$ be a cut of a disk D^2 between $\varphi(v_1)$ and $\varphi(v_2)$, where $\{v_1, v_2\} = V_{ter}$.

Let us fix a homeomorphism

$$\Phi_0 : \partial D^2 \cup \varphi(T) \rightarrow \partial D^2 \cup ([-1, 1] \times \{0\}) ,$$

such that $\Phi_0 \circ \varphi(v_1) = (-1, 0)$, $\Phi_0 \circ \varphi(v_2) = (1, 0)$, $\Phi_0(\partial D^2) = \partial D^2$, $\Phi_0 \circ \varphi(T) = [-1, 1] \times \{0\}$.

By Shernflic's theorem [11, 13] we can find a homeomorphism $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ which extends Φ_0 . It is obvious that an embedding $\Phi \circ \varphi : T \rightarrow \mathbb{R}^2$ complies with the conditions of lemma. From fact that Φ is homeomorphism it follows that φ satisfies conditions of lemma.

Step of induction. Suppose that for some $n > 2$ lemma is proved for all trees with $\sharp V^* < n$ and their embeddings into $\mathbb{R}^2$ which hold Conditions (1) and (2).

Let a tree T such that $V_{ter} \subseteq V^*$, $\sharp V^* = n$, and an embedding $\varphi : T \rightarrow \mathbb{R}^2$ which satisfy Conditions (2) is fixed.

As we noticed above the set V_{ter} contains at least two elements $w_1, w_2 \in V_{ter}$. Let us consider the path $P(w_1, w_2)$ which connects those vertices. Suppose that it passes through the vertices in the following order $w_1 = u_0, u_1, \dots, u_{k-1}, u_k = w_2$. Every vertex $u_1, \dots, u_{k-1}$ has degree at least 2 since it is adjacent to two edges of $P(w_1, w_2)$.

There exists a vertex u_s , $s \in \{1, \dots, k-1\}$ such that

- (i) a degree of u_i equals to 2 and $u_i \notin V^*$ for $i \in \{1, \dots, s-1\}$;
- (ii) either a degree of u_s is greater than 2 or $u_s \in V^*$ and a degree of u_s equals to 2.

Remark that a degree of u_s does not equal to 1. Otherwise, the correlations $u_s = w_2$, $T = P(v_1, v_2)$, $V^* = \{w_1, w_2\}$, $\sharp V^* = 2$ should be satisfied but we assumed that $\sharp V^* \geq 3$.

Let us consider a path $P(w_1, u_s) = P(u_0, u_s)$. Suppose that it passes through edges $e_1, \dots, e_s$ successively.

We consider a subgraph T' of T with the set of vertices and edges, respectively, as follows

$$V(T') = V \setminus \{u_0, \dots, u_{s-1}\}, \quad E(T') = E \setminus \{e_1, \dots, e_s\}.$$

By construction $u_0 \in V_{ter}(T)$ and u_0 is adjacent to e_1 in T ; every vertex u_i , $i \in \{1, \dots, s-1\}$ has degree 2 thus it is adjacent only to e_i and e_{i+1} in T . Therefore a graph T' is defined correctly.

A graph T' has no cycles since it is a subgraph of T . Let us verify that T' is connected. Let $v', v'' \in V(T')$ and $P(v', v'')$ be a path which connects vertices v' and v'' in T . Then a path $P(v', v'')$ does not pass through a vertex $u_0 = w_1$ since $u_0 \in V_{ter}$ and only one edge e_1 is adjacent to this vertex. Thus $e_1 \notin P(v', v'')$. Similarly, if $s \geq 2$ then $e_2 \notin P(v', v'')$ since an edge e_2 is adjacent to a vertex u_1 which is in addition adjacent only to e_1 and $e_1 \notin P(v', v'')$. Similarly, by induction we prove that $e_i \notin P(v', v'')$ for every $i \in \{1, \dots, s\}$. Thus a path $P(v', v'')$ connects vertices v' and v'' in T' . Therefore a graph T' is connected.

We verified that T' is a tree. Let us define $V^*(T') = V^*(T) \cap V(T')$, $\varphi_0 = \varphi|_{T'} : T' \rightarrow \mathbb{R}^2$. By definition of a set $V^*(T')$ it is obvious that a map φ_0 satisfies

condition (2). Also $\#V^*(T') < \#V^*(T)$ since $u_0 \in V^*(T) \setminus V^*(T')$. Thus $\#V^*(T') < n$.

Let us check that $V_{ter}(T') \subseteq V^*(T')$.

By construction for every vertex $v \neq u_s$ of T' its degrees coincide in T and T' . The degree of u_s in T' is one less than degree of u_s in T . Thus $V_{ter}(T') \subseteq V_{ter}(T) \cup \{u_s\}$.

If $u_s \in V^*(T)$, then $V_{ter}(T') \subseteq V_{ter}(T) \cup V^*(T) \subseteq V^*(T)$. Therefore $V_{ter}(T') \subseteq V^*(T) \cap V(T') = V^*(T')$.

Let $u_s \notin V^*(T)$. By definition the degree of u_s in T is not less than 3 and a degree of u_s in T' is not less than 2. Thus $V_{ter}(T') \subseteq V_{ter}(T) \subseteq V^*(T)$. So, as above, $V_{ter}(T') \subseteq V^*(T')$.

By induction lemma holds true for a tree T' and an embedding $\varphi_0 : T' \rightarrow \mathbb{R}^2$.

Denote by $W_0 = \mathbb{R}^2 \setminus D^2, W_1, \dots, W_r$ connected components of a set $\mathbb{R}^2 \setminus (\varphi_0(T') \cup \partial D^2)$.

It is obvious that

$$\varphi(T) = \varphi(T') \cup \varphi(P(u_0, u_s)) = \varphi_0(T') \cup \varphi(P(u_0, u_s)).$$

Therefore $\varphi(T) \cup \partial D^2 = (\varphi_0(T') \cup \partial D^2) \cup \varphi(P(u_0, u_s))$. By construction we get that $(\varphi(T') \cup \partial D^2) \cap \varphi(P(u_0, u_s)) = \{\varphi(u_0), \varphi(u_s)\}$.

Denote $J = \varphi(P(u_0, u_s))$. The set $J_0 = J \setminus \{\varphi(u_0), \varphi(u_s)\}$ is a homeomorphic image of interval thus it is connected. But besides $J_0 \cap (\varphi_0(T') \cup \partial D^2) = \emptyset$ thus there exists a component W_j which contains J_0 (it is easy to see that $j \neq 0$).

By assumption of induction the boundary of disk W_j is a simple closed curve $\partial W_j = K_j \cup \varphi_0(P(v_j, v'_j))$ which consists of an arc K_j of a circle ∂D^2 with the ends $\varphi_0(v_j)$ and $\varphi_0(v'_j)$ and an image of path $P(v_j, v'_j)$ which connects vertices $v_j, v'_j \in V^*(T')$ in T' (this path also connects vertices v_j and v'_j in T).

The set J is a homeomorphic image of segment and also $J_0 \subseteq W_j$, $\varphi(u_0) \in \partial D^2 \subseteq (\mathbb{R}^2 \setminus W_j)$, $\varphi(u_s) \in \varphi_0(T') \subseteq (\mathbb{R}^2 \setminus W_j)$. Therefore J is a cut of disk W_j between points $\varphi(v_j)$ and $\varphi(v'_j)$. Correlations $\varphi(u_s) \in \varphi(P(v_j, v'_j))$, $\varphi(u_0) \in K_j \setminus \{\varphi(v_j), \varphi(v'_j)\} = \partial W_j \setminus \varphi(T')$ hold true since $u_0 \notin V(T')$ and $\varphi(u_0) \notin \varphi(T')$.

So, a set $\overline{W_j} \setminus (\partial W_j \cup \varphi(P(u_0, u_s)))$ has two connected components W_j^1, W_j^2 which are homeomorphic to open disks and bounded by simple closed curves.

We remark that the arc $\varphi(P(v_j, v'_j))$ is not a point, otherwise the correlations $K_j \cong \partial D^2$, $\varphi_0(T') \cap \partial D^2 = \{\varphi(v_j) = \varphi(v'_j)\}$, $\#V^*(T') = \#(\varphi_0(T') \cap \partial D^2) = 1$ should hold true. Thus points $\varphi(v_j)$ and $\varphi(v'_j)$ are different. From the inclusions $\varphi(u_s) \in \varphi(P(v_j, v'_j))$, $\varphi(u_0) \in \partial W_j \setminus \varphi(P(v_j, v'_j))$ it follows that points $\varphi(v_j)$ and $\varphi(v'_j)$ can not be contained in a set $\partial W_j^1 \cap \partial W_j^2 = \varphi(P(u_0, u_s))$ simultaneously.

Let $\varphi(v_j) \in \partial W_j^1$, $\varphi(v'_j) \in \partial W_j^2$. By those correlations the sets W_j^1 and W_j^2 are defined uniquely.

Points $\varphi(u_0), \varphi(u_s)$ divide the circle onto two arcs R_1, R_2 with $R_1 \subseteq \partial W_j^1 \setminus W_j^2$, $R_2 \subseteq \partial W_j^2 \setminus W_j^1$.

Suppose for some edge $e \in E(T)$ its image is contained in ∂W_j . Then the image of e without the ends is connected set and belongs to $\partial W_j \setminus \{\varphi(u_0), \varphi(u_s)\} = R_1 \cup R_2$. Thus the image of e without the endpoints belongs to either R_1 or R_2 .

The path which connects vertices v_j and v'_j in T' passes through the vertices $v_j = \hat{v}_0, \hat{v}_1, \dots, \hat{v}_k = v'_j$ and through the edges $\hat{e}_1, \dots, \hat{e}_k$ in this order.

If $\varphi(\hat{v}_i) \in R_1$ for some $i \in \{0, \dots, k\}$, then $\varphi(\hat{v}_i) \in \mathbb{R}^2 \setminus \overline{R_2}$ and $(\varphi(\hat{e}_i) \setminus \{\varphi(\hat{v}_i), \varphi(\hat{v}_{i+1})\}) \cap (\mathbb{R}^2 \setminus \overline{R_2}) \neq \emptyset$ since a point $\varphi(\hat{v}_i)$ is a boundary for the set $\varphi(\hat{e}_i) \setminus \{\varphi(\hat{v}_i), \varphi(\hat{v}_{i+1})\}$ but $\mathbb{R}^2 \setminus \overline{R_2}$ is an open neighborhood of this point. From what we said it follows that $\varphi(\hat{e}_i) \setminus \varphi(\hat{v}_{i+1}) \subseteq R_1$. Therefore $\varphi(\hat{v}_{i+1}) \in \overline{R_1} = R_1 \cup \{\varphi(u_0), \varphi(u_s)\}$. Indeed, either $\varphi(\hat{v}_{i+1}) \in R_1$ or $\varphi(\hat{v}_{i+1}) = \varphi(u_s)$ (and $\hat{v}_{i+1} = u_s$) since $u_0 \notin V(T')$ by construction.

By assumption of induction $\varphi(u_s) \in \varphi(P(v_j, v'_j)) = \varphi(\hat{v}_0, \hat{v}_k)$. Therefore $u_s \in \{\hat{v}_0, \dots, \hat{v}_k\}$ and there exists an index $k_0 \in \{0, \dots, k\}$ such that $u_s = \hat{v}_{k_0}$.

The inductive application of our previous argument leads us to correlations $\varphi(P(\hat{v}_0, u_s)) \setminus \varphi(u_s) = \varphi(P(v_j, u_s)) \setminus \varphi(u_s) \subseteq R_1$ (in the case when $v_j = u_s$ we get $\varphi(P(v_j, u_s)) = \varphi(u_s)$).

Similar argument give $\varphi(P(u_s, v'_j)) \setminus \varphi(u_s) \subseteq R_2$.

Finally we get $\partial W_j^1 = R_1 \cup \varphi(P(u_0, u_s)) = R_1 \cup J$, $\partial W_j^2 = R_2 \cup J$; $\partial W_j^1 \cap \varphi(P(v_j, v'_j)) = \partial W_j^1 \cap (\varphi(P(v_j, u_s)) \cup \varphi(P(u_s, v'_j))) = \varphi(P(v_j, u_s))$; $\partial W_j^2 \cap \varphi(P(v_j, v'_j)) = \varphi(P(u_s, v'_j))$.

Therefore $\varphi(T) \cap \partial W_j^1 = (\varphi(T') \cup J) \cap \partial W_j^1 = (\varphi(P(v_j, v'_j)) \cup J) \cap \partial W_j^1 = \varphi(P(v_j, u_s)) \cup \varphi(P(u_0, u_s)) = \varphi(P(v_j, u_0))$; $\varphi(T) \cap \partial W_j^2 = \varphi(P(v'_j, u_0))$.

It is easy to see that $\varphi(v_j) \neq \varphi(u_0)$ and $\varphi(v'_j) \neq \varphi(u_0)$ since $v_j, v'_j \in V(T')$ but $u_0 \notin V(T')$. Hence a set $\varphi(P(v_j, u_0)) \setminus \{\varphi(v_j), \varphi(u_0)\}$ is one of two connected components of the set $\partial W_j^1 \setminus \{\varphi(v_j), \varphi(u_0)\}$. Another connected component of this set is contained in $\partial W_j \setminus \varphi(T') = K_j \subseteq \partial D^2$ thus it is an arc of circle ∂D^2 which connects points $\varphi(v_j)$ and $\varphi(u_0)$. Denote it by K_j^1 .

Similarly, $\partial W_j^2 = \varphi(P(v'_j, u_0)) \cup K_j^2$, where K_j^2 is an arc of ∂D^2 which connects points $\varphi(v'_j)$ and $\varphi(u_0)$.

We proved that a compliment $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$ has a finite number of connected components

$$\mathbb{R}^2 \setminus D^2 = W_0, W_1, \dots, W_{j-1}, W_j^1, W_j^2, W_{j+1}, \dots, W_r;$$

and the components W_j^1 and W_j^2 satisfy the conditions of lemma. Finally we remark that the correlations $\partial W_k \cap \varphi(T) = \partial W_k \cap \varphi(T') = \partial W_k \cap \varphi_0(T')$ hold true for $k > 0, k \neq j$ thus

$$\partial W_k = K_k \cup \varphi_0(P(v_k, v'_k)) = K_k \cup \varphi(P(v_k, v'_k))$$

and the component W_k satisfies lemma.

□

Corollary 1.1. *Let T be a tree with fixed subset of vertices $V^* \supseteq V_{ter}$ and $\varphi : T \rightarrow \mathbb{R}^2$ an embedding which satisfies (2).*

Then the following conditions hold true.

1) *In notation of Lemma 1.1*

$$L_i \cap \varphi(T) = \{\varphi(v_i), \varphi(v'_i)\}, \quad i = 1, \dots, m.$$

2) *If there exists an arc L of circle ∂D^2 with the ends $\varphi(u_1), \varphi(u_2)$ such that $L \cap \varphi(T) = \{\varphi(u_1), \varphi(u_2)\}$ for some $u_1, u_2 \in V^*$, then there exists $k \in \{1, \dots, m\}$ such that $L \cup \varphi(P(u_1, u_2)) = \partial U_k$ (then $L = L_k, u_1 = v_k, u_2 = v'_k$).*

Proof. 1) Suppose that an arc $L_i \setminus \{\varphi(v_i), \varphi(v'_i)\}$ contains a point $\varphi(v) \in \varphi(T)$ for some $i \in \{1, \dots, m\}$. Thus $v \in V^*$. Let $e \in E(T)$ be an edge of graph T which is adjacent to a vertex v and $v' \in V$ be another end of the edge e .

A set $J_0 = \varphi(e) \setminus \{\varphi(v), \varphi(v')\}$ is connected, $\varphi(v)$ is a boundary point of it, $W = \mathbb{R}^2 \setminus \varphi(P(v_i, v'_i))$ is an open neighborhood of a point $\varphi(v)$. Thus $J_0 \cap W \neq \emptyset$ and $e \notin P(v_i, v'_i)$. Hence $J_0 \cap \varphi(P(v_i, v'_i)) = \emptyset$. By the conditions of lemma also $J_0 \cap \partial D^2 = \emptyset$. A set $\varphi(P(v_i, v'_i))$ is a cut of closed disk D^2 . Obviously, by construction a set $Q = \overline{U}_i \setminus \varphi(P(v_i, v'_i)) = U_i \cup (L_i \setminus \{\varphi(v_i), \varphi(v'_i)\})$ is a connected component of the compliment $D^2 \setminus \varphi(P(v_i, v'_i))$ which contains a point $\varphi(v)$. That point is a boundary point of the connected subset J_0 of a space $D^2 \setminus \varphi(P(v_i, v'_i))$ therefore $J_0 \subseteq Q$.

But $U_i \subseteq \mathbb{R}^2 \setminus \varphi(T)$, $L_i \subseteq \partial D^2$ and $J_0 \subseteq \varphi(T) \setminus \varphi(V) \subseteq \varphi(T) \setminus \partial D^2$. Thus $J_0 \cap Q \subseteq (J_0 \cap U_i) \cup (J_0 \cap L_i) = \emptyset$.

The contradiction obtained is a last step of the proof of first condition of corollary.

2) Support that there exists an arc L of ∂D^2 with the ends in points $\varphi(u_1), \varphi(u_2)$ such that $L \cap \varphi(T) = \{\varphi(u_1), \varphi(u_2)\}$ for some $u_1, u_2 \in V^*$.

An arc L bounds to some connected component $U_k, k \geq 1$ of the compliment $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$. From Lemma 1.1 and first condition of corollary it follows that $\{u_1, u_2\} = \{v_k, v'_k\}$. Thus vertices u_1 and u_2 can be connected by a path $\tilde{P}(u_1, u_2) = P(v_k, v'_k)$ which satisfies Lemma 1.1. A graph T is a tree thus $P(u_1, u_2) = \tilde{P}(u_1, u_2) = P(v_k, v'_k)$. $\square$

Let T be a tree with a fixed subset of vertices V^* and $\varphi : T \rightarrow \mathbb{R}^2$ is an embedding which satisfy (1) and (2).

The pair of vertices $v_1, v_2 \in V^*, v_1 \neq v_2$ is said to be *adjacent on a circle ∂D^2* if there exists an arc L of this circle with the ends $\varphi(v_1)$ and $\varphi(v_2)$ such that $L \cap \varphi(T) = \{\varphi(v_1), \varphi(v_2)\}$ holds true for it.

Denote by $\mathcal{P}$ a set of all paths in T which connect adjacent pairs of vertices.

Corollary 1.2. *If $\sharp V^* \geq 3$, then a correspondence*

$$\begin{aligned}\Theta : \{U_1, \dots, U_m\} &\rightarrow \mathcal{P}, \\ \Theta(U_i) &= P(v_i, v'_i),\end{aligned}$$

is a bijective map.

Proof. It is sufficient to check an injectivity of the map Θ .

Suppose that the following equalities hold true $\partial U_i = L_i \cup \varphi(P(v, v'))$, $\partial U_j = L_j \cup \varphi(P(v, v'))$ for some $i, j \in \{1, \dots, m\}$, $i \neq j$. Then $L_i \cap L_j = \{\varphi(v), \varphi(v')\}$, $L_i \cup L_j \cong S^1$ therefore $L_i \cup L_j = \partial D^2$.

But from Corollary 1.1 it follows that $\emptyset = (L_i \cup L_j \setminus \{\varphi(v), \varphi(v')\}) \cap \varphi(T)$. Therefore $\sharp(\partial D^2 \cap \varphi(T)) = \sharp V^* \leq 2$ and it contradicts to the conditions of corollary. $\square$

1.2. On relations defined on finite sets. At first we remind that a ternary relation O on the set A is any subset of the 3^{rd} cartesian power A^3 : $O \subseteq A^3$.

Let A be a set, O a ternary relation on A which is asymmetric ($(x, y, z) \in O \Rightarrow (z, y, x) \notin O$), transitive $(x, y, z) \in O, (x, z, u) \in O \Rightarrow (x, y, u) \in O$ and cyclic $(x, y, z) \in O \Rightarrow (y, z, x) \in O$. Then O is called a cyclic order on the set A [12].

A cyclic order O is a complete on a finite set A , $\sharp A \geq 3$, if $x, y, z \in A, x \neq y \neq z \neq x \Rightarrow$ there exists a permutation (u, v, w) of sequence (x, y, z) such that $(u, v, w) \in O$.

Proposition 1.1. *Let there is a complete cyclic order O on some finite set A , $\sharp A \geq 3$.*

Then for every $a \in A$ there exist unique $a', a'' \in A$ such that

- $O(a', a, b)$ for all $b \in A \setminus \{a, a'\}$;
- $O(a, a'', b)$ for all $b \in A \setminus \{a, a''\}$,

and $a' \neq a''$.

Proof. Let us fix $a \in A$. By using [12] we can construct a binary relation ρ up to the relation O with the help of the following condition

$$O(a, a_1, a_2) \Leftrightarrow a_1 \rho a_2.$$

It is easy to verify that the relation ρ defines a strict linear order on a set $A \setminus \{a\}$.

The set $A \setminus \{a\}$ is finite therefore there exist a minimal element a' and maximal element a'' with respect to the order ρ on this set. It is obvious that they satisfy conditions of proposition by definition.

Finally, $a' \neq a''$ since $\sharp(A \setminus \{a\}) \geq 2$. $\square$

Definition 1.1. *Let there is a complete cyclic order O on a set A , $\sharp A \geq 3$. Elements $a_1, a_2 \in A$ are said to be adjacent with respect to a cyclic order O if one of the following conditions holds:*

- $O(a_1, a_2, b)$ for all $b \in A \setminus \{a_1, a_2\}$;

- $O(a_2, a_1, b)$ for all $b \in A \setminus \{a_1, a_2\}$.

Remark 1.1. From Proposition 1.1 it follows that every element has exactly two adjacent elements on a finite set A with a complete cyclic order.

Definition 1.2. Let A be a finite set. A binary relation ρ on A is said to be convenient if

- 1) for all $a, b \in A$ from $a\rho b$ it follows that $a \neq b$;
- 2) for every $a \in A$ there is no more than one $a' \in A$ such that apa' ;
- 3) for every $a \in A$ there is no more than one $a'' \in A$ such that $a''pa$.

We remind that a *graph* of the relation ρ on A is a set $\{(a, b) \in A \times A \mid a\rho b\}$.

Let ρ be a convenient relation on a finite set A , $\hat{\rho}$ be a minimal relation of equivalence which contains ρ . Let us remind that a graph $\hat{\rho}$ consists of

- all pairs (a, b) such that there exist $k = k(a, b) \in \mathbb{N}$ and a sequence $a = a_0, a_1, \dots, a_k = b$ which comply with one of the following conditions $a_{i-1}\rho a_i, a_i\rho a_{i-1}$ for every $i \in \{1, \dots, k\}$;
- pairs $(a, a), a \in A$.

We distinguished a diagonal $\Delta_{A \times A}$ since, in general, there could exist $a \in A$ such that neither $a\rho b$ nor $b\rho a$ holds true for all $b \in A$.

The relation $\hat{\rho}$ generates a partition $\mathfrak{f}$ of A onto classes of equivalence.

Proposition 1.2. Let $B \in \mathfrak{f}$ be a class of equivalence of the relation $\hat{\rho}$. Then there exists no more than one element $b \in B$ which is in the relation ρ with no element of A .

Proof. We remark that if either $a\rho b$ or $b\rho a$ and $b \in B$, then $a \in B$ by definition of B .

It is obvious that if $\sharp B = 1$ then proposition holds true. Let $\sharp B \geq 2$.

Let $a_0, a_1, \dots, a_k$ be a fixed sequence of pairwise different elements of B such that the correlation $a_{i-1}\rho a_i$ holds true for any $i \in \{1, \dots, k\}$.

If there exists $b \in B \setminus \{a_0, \dots, a_k\}$, then there exists $b' \in B \setminus \{a_0, \dots, a_k\}$ such that either $b'\rho a_0$ or $a_k\rho b'$. Let us verify it.

By definition of a set B there exists a sequence $b = c_0, c_1, \dots, c_m = a_0$ such that either $c_{j-1}\rho c_j$ or $c_j\rho c_{j-1}$ holds true for all $j \in \{1, \dots, m\}$. From correlations $c_0 \notin \{a_0, \dots, a_k\}$ and $c_m \in \{a_0, \dots, a_k\}$ it follows that there is $s \in \{0, \dots, m\}$ such that $c_{s-1} \notin \{a_0, \dots, a_k\}$ but $c_s \in \{a_0, \dots, a_k\}$. Thus $c_s = a_r$ for some $r \in \{0, \dots, k\}$.

Let $c_{s-1}\rho c_s$, i.e. $c_{s-1}\rho a_r$. Then $r = 0$. Really, if $r \geq 1$, then $a_{r-1}\rho a_r$. By construction $c_{r-1} \neq a_{r-1}$ therefore a correlation $c_{s-1}\rho a_r$ contradicts to condition 3) of Definition 1.2.

Similarly, if $c_s\rho c_{s-1}$, then $c_s = a_k$.

It is easy to see that element $b' = c_{s-1}$ satisfies conditions of proposition.

From what we said above it follows that if for some pairwise different $a_0, \dots, a_k \in B$ inequality $\{a_0, \dots, a_k\} \neq B$ and relation $a_{i-1}\rho a_i$, $i \in \{1, \dots, k\}$ hold true, then there are pairwise different $a'_0, \dots, a'_{k+1} \in B$ such that $a'_{i-1}\rho a'_i$, $i \in \{1, \dots, k+1\}$ hold true for them.

By definition the set B contains two elements $b', b'' \in B$ such that $b'\rho b''$. So, by a finite number of steps (the set B is finite) we can index all elements of B in such way that the following correlations hold true

$$(3) \quad \begin{aligned} a_{i-1}\rho a_i, \quad i \in \{1, \dots, n\}; \\ \{a_0, \dots, a_n\} = B. \end{aligned}$$

Therefore only element $a_n \in B$ can satisfy conditions of the proposition. $\square$

Let μ be some relation on a set A .

Definition 1.3. Elements $b_0, \dots, b_n \in A$, $n \geq 1$ are said to generate μ -cycle if a graph of the relation μ contains a set

$$(4) \quad \{(b_0, b_1), \dots, (b_{n-1}, b_n), (b_n, b_0)\}.$$

Definition 1.4. Elements $b_0, \dots, b_n \in A$, $n \geq 0$ to generate μ -chain if for arbitrary $a \in A$ the pairs (a, b_0) and (b_n, a) do not belong to a graph of μ and for $n \geq 1$ a graph of the relation μ contains a set

$$(5) \quad \{(b_0, b_1), \dots, (b_{n-1}, b_n)\}.$$

Corollary 1.3. Let ρ be a convenient relation, $B \in \mathfrak{f}$ a class of equivalence of the relation $\hat{\rho}$. Then the elements of B generate either ρ -cycle or ρ -chain. In the first case a graph of the restriction of ρ on the set B is of form (4) and in the other it has form (5).

Proof. Let us order the elements of B in such way that (3) holds true for them.

If there exists $a \in A$ such that $a_n\rho a$, then $a \in B$. Conditions 1) and 3) of Definition 1.2 obstruct to hold correlation $a_i\rho a_j$ for $i \neq j-1$, $j \in \{1, \dots, n\}$. Therefore $a = a_0$ and $a_n\rho a_0$.

Similarly, if there exists $a \in A$ which $a\rho a_0$, then from conditions 1) and 2) of Definition 1.2 it follows that $a = a_n$ and $a_n\rho a_0$.

So, either a correlation $a_n\rho a_0$ holds true or for every $a \in A$ neither $a\rho a_0$ nor $a_n\rho a$ holds true. In the first case the elements of B generate ρ -cycle (if $a_n\rho a_0$, then $a_n \neq a_0$ and $\sharp B \geq 2$ by definition), in the other case we get ρ -chain. $\square$

Corollary 1.4. Let the elements of $B \subseteq A$ generate either ρ -cycle or ρ -chain. Then B is a class of equivalence of the relation $\hat{\rho}$. If the elements of $B \subseteq A$ generate ρ -chain, then the relation ρ generates a full linear order on B .

Proof. Let $\hat{\rho}$ be a minimal relation of equivalence which contains ρ . By definition the set B belongs to the unique class of equivalence of the relation $\hat{\rho}$. Denote it by $\hat{B}$.

By definition the set B satisfies (3). If there exists $b \in \hat{B} \setminus B$, then, as we verified in the proof of Proposition 1.2, there is $b' \in \hat{B} \setminus B$ such that

$$(6) \quad b' \rho a_0 \quad \text{or} \quad a_n \rho b'.$$

This contradicts to definition of ρ -chain. If the elements of B generate ρ -cycle, then it follows from the definition of convenient relation that

$$(7) \quad a_n \rho a_0,$$

see Corollary 1.3. By using conditions 2) and 3) of a convenient relation from equality $b' \notin B$ we can conclude that (6) and (7) can not be satisfied simultaneously.

So, a set B is a class of equivalence of the relation $\hat{\rho}$.

If elements of the set B generate a chain, then a graph of a restriction of the relation ρ on B has form (5), see Corollary 1.3. Therefore ρ generates a linear order on the set B . $\square$

Definition 1.5. Let O be a complete cyclic order on A , $\sharp A \geq 3$. O is said to induce a binary relation ρ_O on a A according to the following rule: $a \rho_O b$ if $O(a, b, c) \forall c \in A \setminus \{a, b\}$.

From Proposition 1.1 it follows that a relation ρ_O is convenient.

Proposition 1.3. If O is a complete cyclic order on A , then all elements of A generate ρ_O -cycle.

Proof. Let $\hat{\rho}_O$ be a minimal relation of equivalence which contains ρ_O . From Proposition 1.1 and Corollaries 1.3 and 1.4 it follows that every class of equivalence of the relation $\hat{\rho}_O$ is ρ_O -cycle and there are no any other ρ_O -cycles.

Let $B = \{b_0, \dots, b_k\}$ be some class of equivalence of the relation $\hat{\rho}_O$ and the following correlations are satisfied

$$b_0 \rho b_1, \dots, b_{k-1} \rho b_k, b_k \rho b_0.$$

Support that $B \subsetneq A$. Let us fix $a \in A \setminus B$. By definition the following correlations hold true

$$\begin{aligned} O(b_{i-1}, b_i, a), \quad i \in \{1, \dots, k\}; \\ O(b_k, b_0, a). \end{aligned}$$

Thus it follows from definition of cyclic order it follows that

$$\begin{aligned} O(a, b_{i-1}, b_i), \quad i \in \{1, \dots, k\}; \\ O(a, b_k, b_0). \end{aligned}$$

From definition it also follows that if $O(a, b_0, b_{i-1})$ and $O(a, b_{i-1}, b_i)$, then $O(a, b_0, b_i)$. Therefore starting from $O(a, b_0, b_1)$ in the finite number of steps we get $O(a, b_0, b_k)$.

Thus $O(a, b_k, b_0)$ and $O(a, b_0, b_k)$ should be satisfied simultaneously but it contradicts to antisymmetry of cyclic order.

Therefore all elements of a set A are equivalent under $\hat{\rho}_O$ and generate ρ_O -cycle. $\square$

Definition 1.6. Let ρ be a convenient relation on a finite set A . We define a ternary relation O_ρ on A with the help of the following rule. The ordered triple (a_1, a_2, a_3) of A is said to be in the relation O_ρ if $a_1 \neq a_2 \neq a_3 \neq a_1$ and there are

$$(8) \quad a_1 = a_0^{12}, a_1^{12}, \dots, a_{m(1)}^{12} = a_2 = a_0^{23}, \dots, a_{m(2)}^{23} = a_3 = a_0^{31}, \dots, a_{m(3)}^{31} = a_1,$$

which satisfy the following conditions:

- $a_{n-1}^{sr} \rho a_n^{sr}$ for all $n \in \{1, \dots, m(s)\}$ and $(s+1) \equiv r \pmod{3}$;
- $a_n^{sr} \notin \{a_1, a_2, a_3\}$ for all $n \in \{1, \dots, m(s)-1\}$ and $(s+1) \equiv r \pmod{3}$.

Proposition 1.4. The relation O_ρ is a cyclic order on A .

Proof. From definition it is obvious that the relation O_ρ is cyclic.

Let us remark that from definition if $O_\rho(a_1, a_2, a_3)$, then all elements of a set (8) (in particular elements a_1, a_2 and a_3) belong to the same class of equivalence of minimal equivalence relation $\hat{\rho}$ which contains ρ .

We should verify that all elements a_n^{sr} , $n \in \{1, \dots, m(s)\}$, $(s+1) \equiv r \pmod{3}$ are different.

Suppose that it is not true and there are two different sets of indexes such that $a_n^{sr} = a_k^{t\tau}$, $n \in \{1, \dots, m(s)\}$, $k \in \{1, \dots, m(t)\}$, $(s+1) \equiv r \pmod{3}$, $(t+1) \equiv \tau \pmod{3}$.

Let us consider two sequences

$$\begin{aligned} (b_1, \dots, b_i) &= (a_n^{sr}, a_{n+1}^{sr}, \dots, a_{m(s)}^{sr}, \dots, a_0^{t\tau}, a_1^{t\tau}, \dots, a_{k-1}^{t\tau}, a_k^{t\tau}), \\ (c_1, \dots, c_j) &= (a_k^{t\tau}, a_{k+1}^{t\tau}, \dots, a_{m(t)}^{t\tau}, \dots, a_0^{sr}, a_1^{sr}, \dots, a_{n-1}^{sr}, a_n^{sr}). \end{aligned}$$

Those two sequences satisfy the following conditions:

- $b_{l-1} \rho b_l$ for all $l \in \{1, \dots, i\}$;
- $c_{l-1} \rho c_l$ for all $l \in \{1, \dots, j\}$;
- $b_i = c_1 = c_j = b_1$;
- there exists $\hat{a} \in \{a_1, a_2, a_3\}$ such that either $\hat{a} \in \{b_1, \dots, b_i\} \setminus \{c_1, \dots, c_j\}$ or $\hat{a} \in \{c_1, \dots, c_j\} \setminus \{b_1, \dots, b_i\}$ since every element a_1, a_2, a_3 is contained exactly once in the sequence (8) by definition.

Let $\hat{a} \notin \{b_1, \dots, b_i\}$. By definition the elements $b_1, \dots, b_i$ generate a cycle therefore the set $\{b_1, \dots, b_i\}$ is a class of equivalence of the relation $\hat{\rho}$, see Corollary 1.4. But it contradicts to the condition that all elements of the set (8) belong to the same class of equivalence of the relation $\hat{\rho}$.

The case when $\hat{a} \notin \{c_1, \dots, c_j\}$ can be considered similarly.

Therefore all elements of the set (8) are different.

Let $O_\rho(a_1, a_2, a_3)$ and $O_\rho(a_3, a_2, a_1)$ hold true simultaneously. Then from definition there are two sequences $a_3 = a_0^{31}, a_1^{31}, \dots, a_{m(3)}^{31} = a_1$ and $a_1 = b_0^{31}, b_1^{31}, \dots, b_{n(3)}^{31} = a_3$ such that

- $a_{i-1}^{31} \rho a_i^{31}$ for all $i \in \{1, \dots, m(3)\}$;
- $b_{j-1}^{31} \rho b_j^{31}$ for all $j \in \{1, \dots, n(3)\}$;
- $a_2 \notin \{a_0^{31}, \dots, a_{m(3)}^{31}, b_0^{31}, \dots, b_{n(3)}^{31}\}$.

It is obvious that there is $k \in \{1, \dots, m(3)\}$ such that $a_i^{31} \notin \{b_0^{31}, \dots, b_{n(3)}^{31}\}$ for $i < k$ but $a_k^{31} \in \{b_0^{31}, \dots, b_{n(3)}^{31}\}$. Hence $a_k^{31} = b_l^{31}$ for some $l \in \{1, \dots, n(3)\}$ and $a_k^{31} \rho b_{l+1}^{31}$. It is clear that all elements of the following sequence

$$a_3 = a_0^{31}, \dots, a_k^{31}, b_{l+1}^{31}, \dots, b_{n(3)}^{31}$$

are different and generate ρ -cycle. Further by definition a_2 does not belong to that sequence. Therefore $a_3 = a_0^{31}$ and a_2 belong to different classes of equivalence of relation $\hat{\rho}$, see Corollary 1.4.

On the other hand elements a_1, a_2 and a_3 must belong to the unique class of equivalence $\hat{\rho}$, see above.

This contradiction proves the antisymmetry of the relation O_ρ .

Let $O_\rho(a_1, a_2, a_3)$ and $O_\rho(a_1, a_3, a_4)$ for some $a_1, \dots, a_4 \in A$.

We should remark that the elements $a_1, \dots, a_4$ are pairwise different. Really, by definition $a_1 \neq a_3$ and $\{a_1, a_3\} \cap \{a_2, a_4\} = \emptyset$. If $a_2 = a_4$, then from a cyclicity of relation O_ρ it follows that $O_\rho(a_3, a_1, a_2)$ and $O_\rho(a_4, a_1, a_3) = O_\rho(a_2, a_1, a_3)$. But it is impossible since a relation O_ρ is antisymmetric.

Let us consider a sequence (8). Its elements generate ρ -cycle. We will prove that $a_4 \in \{a_1^{31}, \dots, a_{m(3)-1}^{31}\}$.

Suppose that $a_4 \in \{a_1^{12}, \dots, a_{m(1)-1}^{12}\}$. Then $a_4 = a_k^{12}$, $k \in \{1, \dots, m(1) - 1\}$. We consider the sequences

$$\begin{aligned} (b_0^{12}, \dots, b_{t(1)}^{12}) &= (a_1 = a_0^{12}, \dots, a_k^{12} = a_4); \\ (b_0^{23}, \dots, b_{t(2)}^{23}) &= (a_4 = a_k^{12}, \dots, a_{m(1)}^{12} = a_0^{23}, \dots, a_{m(2)}^{23} = a_3); \\ (b_0^{31}, \dots, b_{t(3)}^{31}) &= (a_3 = a_0^{31}, \dots, a_{m(3)}^{31} = a_1). \end{aligned}$$

Join them into a sequence

$$a_1 = b_0^{12}, \dots, b_{t(1)}^{12} = a_4 = b_0^{23}, \dots, b_{t(2)}^{23} = a_3 = b_0^{31}, \dots, b_{t(3)}^{31}.$$

By the construction all elements of such sequence generate ρ -cycle therefore it satisfies the properties which are similar to the conditions of the sequence (8). We get $O_\rho(a_1, a_4, a_3)$. Then from a cyclicity of the relation O_ρ it follows that $O_\rho(a_4, a_3, a_1)$. But by the condition we have $O_\rho(a_1, a_3, a_4)$, moreover, we proved that the relation O_ρ is antisymmetric. Thus the relation $O_\rho(a_4, a_3, a_1)$ does not hold true and $a_4 \notin \{a_1^{12}, \dots, a_{m(1)-1}^{12}\}$.

The fact that $a_4 \notin \{a_1^{23}, \dots, a_{m(2)-1}^{23}\}$ can be proved similarly.

Therefore $a_4 \in \{a_1^{31}, \dots, a_{m(3)-1}^{31}\}$ and $a_4 = a_s^{31}$ for some $s \in \{1, \dots, m(3) - 1\}$.

Let us consider the sequences

$$\begin{aligned}(c_0^{12}, \dots, c_{\tau(1)}^{12}) &= (a_1 = a_0^{12}, \dots, a_{m(1)}^{12} = a_2); \\ (c_0^{23}, \dots, c_{\tau(2)}^{23}) &= (a_2 = a_0^{23}, \dots, a_{m(2)}^{23} = a_0^{31}, \dots, a_s^{31} = a_4); \\ (c_0^{31}, \dots, c_{\tau(3)}^{31}) &= (a_4 = a_s^{31}, \dots, a_{m(3)}^{31} = a_1).\end{aligned}$$

Let us join them into a sequence

$$a_1 = c_0^{12}, \dots, c_{\tau(1)}^{12} = a_2 = c_0^{23}, \dots, c_{\tau(2)}^{23} = a_4 = c_0^{31}, \dots, c_{\tau(3)}^{31}.$$

By construction this sequence satisfies the conditions of definition 1.6. Therefore the correlation $O_\rho(a_1, a_2, a_4)$ holds true and the relation O_ρ is transitive.

Finally, we can conclude that the relation O_ρ satisfies all conditions of definition of cyclic order. $\square$

Definition 1.7. Let C and D be cyclic orders on sets A and B , respectively. Let $\varphi : A \rightarrow B$ be a bijective map.

A map φ is called a monomorphism of cyclic order C into a cyclic order D if $C(a_1, a_2, a_3) \Rightarrow D(\varphi(a_1), \varphi(a_2), \varphi(a_3))$; it is called an epimorphism C onto D if $D(b_1, b_2, b_3) \Rightarrow C(\varphi^{-1}(b_1), \varphi^{-1}(b_2), \varphi^{-1}(b_3))$; φ is an isomorphism C onto D if $C(a_1, a_2, a_3) \Leftrightarrow D(\varphi(a_1), \varphi(a_2), \varphi(a_3))$.

Remark 1.2. It is clear that

- 1) if φ is a monomorphism of cyclic order C onto D , then φ^{-1} is an epimorphism of D onto C ;
- 2) an isomorphism of the relations of cyclic order is a map which is a monomorphism and an epimorphism simultaneously;
- 3) a relation of isomorphism is a relation of equivalence.

Lemma 1.2. Let C and D be complete cyclic orders on the sets A and B , respectively, $\varphi : A \rightarrow B$ is a bijective map.

If φ is either monomorphism or an epimorphism, then φ is an isomorphism.

Proof. Let φ be an epimorphism (in the case when φ is a monomorphism we consider a map φ^{-1}). Let us check that φ is also a monomorphism.

Let $C(a_1, a_2, a_3)$ for some $a_1, a_2, a_3 \in A$. We define $b_i = \varphi(a_i) \in B$, $i = 1, 2, 3$. From definition it follows that $a_1 \neq a_2 \neq a_3 \neq a_1$. Then $b_1 \neq b_2 \neq b_3 \neq b_1$.

The cyclic order D is full therefore there is a permutation $\sigma \in S(3)$ such that $D(b_{\sigma(1)}, b_{\sigma(2)}, b_{\sigma(3)})$. From an epimorphism of φ we can conclude that $C(a_{\sigma(1)}, a_{\sigma(2)}, a_{\sigma(3)})$. Thus σ is even permutation. Now from antisymmetry and cyclicity of D it follows that $D(b_1, b_2, b_3)$, see [12]. Therefore we get $D(\varphi(a_1), \varphi(a_2), \varphi(a_3))$ and φ is a monomorphism. $\square$

Remark 1.3. Lemma 1.2 holds true for arbitrary sets A and B , i.e. they can be infinite.

Lemma 1.3. *Let O is a relation of complete cyclic order on the finite set A . Then*

$$O = O_{\rho_O},$$

where ρ_O is a convenient binary relation generated by O and O_{ρ_O} is a relation of cyclic order generated by the convenient relation ρ_O .

Proof. We should prove that the relation O_{ρ_O} is full.

Let b_1, b_2, b_3 be some pairwise different elements of A . From Proposition 1.3 and Corollary 1.4 the minimal relation of equivalence $\hat{\rho}_O$ which contains ρ_O has the unique class of equivalence $B = A$. Thus we can index all elements of A in such way that (3) holds true. From Corollary 1.3 we also get $a_n \rho_O a_0$.

It is obvious that $\{b_1, b_2, b_3\} = \{a_{k_1}, a_{k_2}, a_{k_3}\}$ for some $0 \leq k_1 < k_2 < k_3 \leq n$ further there is a inversion $\sigma \in S(3)$ such that $a_{k_i} = b_{\sigma(i)}$, $i = 1, 2, 3$.

Let us consider the sequences

$$\begin{aligned} (c_0^{12}, \dots, c_{m(1)}^{12}) &= (a_{k_1}, a_{k_1+1}, \dots, a_{k_2}); \\ (c_0^{23}, \dots, c_{m(2)}^{23}) &= (a_{k_2}, \dots, a_{k_3}); \\ (c_0^{31}, \dots, c_{m(3)}^{31}) &= (a_{k_3}, \dots, a_n, a_0, \dots, a_{k_1}). \end{aligned}$$

We can join them into one

$$a_{k_1} = c_0^{12}, \dots, c_{m(1)}^{12} = a_{k_2} = c_0^{23}, \dots, c_{m(2)}^{23} = a_{k_3} = c_0^{31}, \dots, c_{m(3)}^{31} = a_{k_1}.$$

By construction this sequence satisfies the conditions of Definition 1.6 thus we get $O_{\rho_O}(a_{k_1}, a_{k_2}, a_{k_3})$. It means that $O_{\rho_O}(b_{\sigma(1)}, b_{\sigma(2)}, b_{\sigma(3)})$ and O_{ρ_O} is full.

Suppose that $O_{\rho_O}(a_1, a_2, a_3)$ holds true for some $a_1, a_2, a_3 \in A$. From Definition 1.6 it follows that there is a sequence

$$a_1 = a_0^{12}, \dots, a_{m(1)}^{12} = a_2,$$

such that $a_{i-1}^{12} \rho_O a_i^{12}$ for all $i \in \{1, \dots, m(1)\}$. Therefore from definition of the relation ρ_O correlations $O(a_{i-1}^{12}, a_i^{12}, a)$ follow for all $a \in A \setminus \{a_{i-1}^{12}, a_i^{12}\}$, $i \in \{1, \dots, m(1)\}$. In particular, $O(a_{i-1}^{12}, a_i^{12}, a_3)$, $i \in \{1, \dots, m(1)\}$. From cyclicity of O it follows that the correlations $O(a_3, a_{i-1}^{12}, a_i^{12})$, $i \in \{1, \dots, m(1)\}$ hold true.

Starting from the correlation $O(a_3, a_0^{12}, a_1^{12}) = O(a_3, a_1, a_1^{12})$, using the previous correlations and transitivity of O we inductively get that $O(a_3, a_1, a_i^{12})$, $i \in \{1, \dots, m(1)\}$. In particular, $O(a_3, a_1, a_{m(1)}^{12}) = O(a_3, a_1, a_2)$. From a cyclicity of O it follows that $O(a_1, a_2, a_3)$.

Therefore an identical map $Id_A : A \rightarrow A$ induces an epimorphism of a complete cyclic order O onto a complete cyclic order O_{ρ_O} . From Lemma 1.2 it follows that the map Id_A is an isomorphism of the cyclic orders O and O_{ρ_O} therefore $O = O_{\rho_O}$. $\square$

Lemma 1.4. *Let ρ be a convenient relation such that all elements of a set A , $\sharp A \geq 3$ generate a cycle.*

Suppose that a graph of relation μ on A is obtained from a graph of ρ by throwing out two pairs (b_1, b'_1) and (b_2, b'_2) (the cases when either $b'_1 = b_2$ or $b'_2 = b_1$ are included). Let $\hat{\mu}$ be a minimal relation of equivalence which contains μ .

Then the relation μ is convenient, $\hat{\mu}$ has exactly two classes of equivalence B_1 and B_2 such that the elements of each of them generate μ -chain and the elements $b_1, b_2 \in A$ belong to the different classes of equivalence of $\hat{\mu}$.

Proof. The fact that μ is a convenient relation is trivial corollary from definition.

The relation μ does not contain cycles. In fact, if the elements of some set $B \subseteq A$ generate μ -cycle, then elements of B generate ρ -cycle. From Corollary 1.4 and the condition of lemma we get $B = A$. Then from definition of a cycle it follows that there is $a \in A$ such that $b_1 \mu a$, hence $b_1 \rho a$. But $b_1 \rho b'_1$ and $b'_1 \neq a$ (by condition of lemma b_1 is not in the relation μ with b'_1). It contradicts to the Condition 2) of definition 1.2.

Thus every class of equivalence of the relation $\hat{\mu}$ is a chain, see Corollary 1.3, and it contains exactly one element which is in the relation μ with no element of A .

By condition of lemma the elements of A generate ρ -cycle. From Definition 1.2 it follows that there is the unique $a' \in A$ such that $a \rho a'$ for every $a \in A$. Then $a \mu a'$, if $a \notin \{b_1, b_2\}$ but b_1 and b_2 are the unique elements of the set A which are not in the relation μ with any element of A .

now the statement of lemma elementary follows from what we said before. $\square$

1.3. A local connectivity of two dimensional disk in boundary points.

Definition 1.8. [11, 13] Let E be a subset of a topological space S and x is some point of S (x does not necessarily belong to E). A set E is called a locally connected in a point x if for every neighborhood U of x there is a neighborhood $U' \subseteq U$ of x such that any two points which belong to $U' \cap E$ can be joined by a connected set which belongs to $U \cap E$.

Lemma 1.5. Let D^2 be a closed two dimensional disk, $x \in \partial D^2$ and W an open neighborhood of point x in a space D^2 .

If for some connected components W_1 and W_2 of a set $W \cap (D^2 \setminus \partial D^2)$ the following correlation holds true $x \in \overline{W_1} \cap \overline{W_2}$, then $W_1 = W_2$.

Proof. Obviously, we can assume that D^2 is a standard two dimensional disk on a plane. Let U be a neighborhood of point x in $\mathbb{R}^2$ such that $D^2 \cap U = W$. It is known, see [11, 13], that every Jordan domain on the plane is locally connected in all points of its boundary. Therefore there exists a neighborhood U' of x such that arbitrary two points which belong to $U' \cap (D^2 \setminus \partial D^2)$ can be connected by a connected set that is contained in $U \cap (D^2 \setminus \partial D^2)$. Therefore all points of the set $U' \cap (D^2 \setminus \partial D^2)$ should belong to the unique connected component of a set $W \cap (D^2 \setminus \partial D^2)$. $\square$

2. CRITERION OF A $\mathcal{D}$ -PLANARITY OF A TREE.

Let T be a tree, V a set of its vertices, V_{ter} a set of its terminal vertices and $V^* \subseteq V$ a subset of T such that $V_{ter} \subseteq V^*$. We assume that if $\#V^* \geq 3$ then there is some cyclic order C defined on V^* .

Let

$$D^2 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$$

be a closed oriented 2-dimensional disk.

Definition 2.1. A tree T is called $\mathcal{D}$ -planar if there exists an embedding $\varphi : T \rightarrow \mathbb{R}^2$ which satisfies (2) and if $\#V^* \geq 3$ then a cyclic order $\varphi(C)$ on $\varphi(V^*)$ coincides with a cyclic order which is generated by the orientation of $\partial D^2 \cong S^1$.

Remark 2.1. A map $\varphi|_{V^*} : V^* \rightarrow \varphi(V^*)$ is bijective whence a ternary relation $\varphi(C)$ on $\varphi(V^*)$ defined by following correlation

$$C(v_1, v_2, v_3) \Rightarrow \varphi(C)(\varphi(v_1), \varphi(v_2), \varphi(v_3)), \quad v_1, v_2, v_3 \in V^*,$$

is a relation of cyclic order.

Remark 2.2. We can define a cyclic order in a natural way on an oriented circle S^1 : an ordered triple of points $x_1, x_2, x_3 \in S^1$ is cyclically ordered if these points are passed in that order in the process of moving along a circle in a positive direction.




FIGURE 1. On the left a tree is $\mathcal{D}$ -planar.

Theorem 2.1. If V^* contains just two vertices, a tree T is $\mathcal{D}$ -planar.

If $\#V^* \geq 3$ then a $\mathcal{D}$ -planarity of T is equivalent to satisfying the following condition:

- for any edge e there are exactly two paths such that they pass through an edge e and connect two adjacent vertices of V^* .

Proof. If $\#V^* = 2$, then T is homeomorphic to a segment and a set of its terminal vertices coincides with $V^* = V_{ter}$, see Lemma 1.1. It is obvious that there exists an embedding $\varphi : T \rightarrow \mathbb{R}^2$ satisfying Definition 2.1 and a tree T is $\mathcal{D}$ -planar.

Let $\#V^* \geq 3$ and T is $\mathcal{D}$ -planar. It means that there is an embedding $\varphi : T \rightarrow \mathbb{R}^2$ which satisfies Definition 2.1.

Let $e \in E(T)$ be an edge of T connecting vertices $w_1, w_2 \in V$. We fix a point $x \in \varphi(e) \setminus \{\varphi(w_1), \varphi(w_2)\}$.

A topological space T is one-dimensional compact hence its homeomorphic image $\varphi(T)$ is one-dimensional [1]. Then $x \in \overline{(\mathbb{R}^2 \setminus \varphi(T))}$. It follows from (2) that $x \in \text{Int } D^2$, therefore

$$x \in \overline{(\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2))}.$$

By Lemma 1.1 there is a connected component U_j of a set $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$ such that a point x belongs to a boundary of it.

Corollary 1.1 states that $\partial U_j \cap \varphi(T) = \varphi(P(v_j, v'_j))$, where $\varphi(v_j), \varphi(v'_j)$ are adjacent with respect to a cyclic order of $\varphi(V^*)$ induced from ∂D^2 , see Remark 2.2. According to Definition 2.1, it is the same as vertices v_j and v'_j are adjacent under a cyclic order C on V^* .

So $e \in P(v_j, v'_j)$ and vertices v_j, v'_j are adjacent. It means that for any edge of a $\mathcal{D}$ -planar tree T there is at least one path that satisfies the condition of theorem.

There exists an open neighborhood $W = e \setminus \{w_1, w_2\}$ of a point $\varphi^{-1}(x)$ in T that is homeomorphic to an interval. Using the compactness of $T \setminus W$ and theorem of Shenflies [11, 13] we can find a neighborhood U of x in $\mathbb{R}^2 \setminus \partial D^2$ and a homeomorphism $h : U \rightarrow \text{Int } D^2$ such that $h(x) = (0, 0)$, $h \circ \varphi(T) = h \circ \varphi(W) = (-1, 1) \times \{0\}$. Let us designate

$$\begin{aligned} U^+ &= h^{-1}(\{(x, y) \in \text{Int } D^2 \mid y > 0\}), \\ U^- &= h^{-1}(\{(x, y) \in \text{Int } D^2 \mid y < 0\}). \end{aligned}$$

It is clear that $U \subseteq \varphi(T) \cup U^+ \cup U^-$. If for some component U_k of $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$ the intersections $U^+ \cap U_k$ and $U^- \cap U_k$ are empty, then $x \notin \overline{U_k}$ and $e \notin P(v_k, v'_k)$ in terms of Lemma 1.1.

By the construction, the sets U^+ and U^- are connected and they belong to $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$. Thus there are two components U_i and U_j such that $U^+ \subseteq U_i$, $U^- \subseteq U_j$, $x \in \overline{U_i} \cap \overline{U_j}$ and $e \in P(v_i, v'_i) \cap P(v_j, v'_j)$.

By Corollaries 1.1 and 1.2 for any edge of T there are no more than two paths such that they connect adjacent vertices of V^* .

In order to verify that there are exactly two such paths it is sufficient to prove that $U_i \neq U_j$.

Suppose that for some component U_i of $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$ we get $U \setminus \varphi(T) = U^+ \cup U^- \subseteq U_i$. An open connected subset U_i of $\mathbb{R}^2$ is path-connected [2].

Denote $a_0^+ = (0, 1/2)$, $a_0^- = (0, -1/2) \in \text{Int } D^2$, $\gamma_0 = \{0\} \times [-1/2, 1/2] \subseteq \text{Int } D^2$, $a^+ = h^{-1}(a_0^+)$, $a^- = h^{-1}(a_0^-) \in U$, $\gamma = h^{-1}(\gamma_0)$.

It is obvious that the points a_0^+ and a_0^- are attainable from domains $h(U^+) \setminus \gamma_0$ and $h(U^-) \setminus \gamma_0$ by a simple continuous curve. Therefore the points a^+ and a^- are attainable from the domain $U_i \setminus \gamma$ and there is a cut $\hat{\gamma}$ of $U_i \setminus \gamma$ between a^+ and a^- [11, 13].

Then $\mu = \gamma \cup \hat{\gamma}$ is a simple close curve such that $\mu \cap \varphi(T) = \{x\}$, $\mu \setminus \{x\} \subseteq U_i$ and $h(\mu) \supseteq \gamma_0$.

By Jordan's theorem μ bounds an open disk G [11, 13].

The point x does not belong to the compact $\hat{\gamma}$ hence there exists its open neighborhood $\hat{U} \subseteq U$ such that $\hat{U} \cap \hat{\gamma} = \emptyset$. Since h maps a neighborhood $\hat{U}$ of a point x into an open neighborhood of origin then there exists an $\varepsilon \in (0, 1/2)$ such that a set

$$Q_0 = \{(x, y) \in D^2 \mid x^2 + y^2 < \varepsilon^2\}$$

does not intersect the set $h(\hat{\gamma})$. It follows that

$$Q_0 \cap h(\varphi(T) \cup \partial D^2) = Q_0 \cap h \circ \varphi(e) = (-\varepsilon, \varepsilon) \times \{0\},$$

$$Q_0 \cap h(\mu) = Q_0 \cap \gamma_0 = \{0\} \times (-\varepsilon, \varepsilon).$$

Denote $Q = h^{-1}(Q_0)$. Evidently, a set Q is an open neighborhood of x .

Open sets

$$h^{-1}(\{(x, y) \in Q_0 \mid x < 0\}) \text{ and } h^{-1}(\{(x, y) \in Q_0 \mid x > 0\})$$

are connected and do not intersect the set μ . Therefore one of them must be contained in a disk G , another should belong to an unbounded domain $\mathbb{R}^2 \setminus \overline{G}$.

Sets $h^{-1}((-\varepsilon, 0) \times \{0\})$ and $h^{-1}((0, \varepsilon) \times \{0\})$ belong to the intersection of these domains with the image $\varphi(e)$ of e . Hence $\varphi(e) \cap G \neq \emptyset$ and $\varphi(e) \cap \mathbb{R}^2 \setminus \overline{G} \neq \emptyset$ hold true.

A segment $\varphi(e)$ is divided by x on two connected arcs that have no common points with $\mu = \partial G$. Thus one of them should belong to G and the other is contained in $\mathbb{R}^2 \setminus \overline{G}$.

Finally, the following statement is true: either $\varphi(w_1)$ or $\varphi(w_2)$ belongs to G and the other point is contained in $\mathbb{R}^2 \setminus \overline{G}$.

Let $\varphi(w_1) \in G$, $\varphi(w_2) \in \mathbb{R}^2 \setminus \overline{G}$.

By the construction, curves ∂D^2 and μ have no common points since either $G \subseteq \text{Int } D^2$ or $\text{Int } D^2 \subseteq G$. But $\emptyset \neq (\gamma \cap Q) \subseteq (\mu \cap U_i) \subseteq (\mu \cap \text{Int } D^2)$. Therefore $\overline{G} \subseteq \text{Int } D^2$.

Let us denote by $\hat{T}$ a graph with a set of vertices $V(\hat{T}) = V(T) = V$ and a set of edges $E(\hat{T}) = E(T) \setminus \{e\} = E \setminus \{e\}$.

It is easy to show that the graph $\hat{T}$ has two connected components $T_1 \ni w_1$ and $T_2 \ni w_2$. The images of them do not intersect with the curve μ , therefore a set $\varphi(T_1)$ together with the point $\varphi(w_1)$ belongs to $G \subseteq \text{Int } D^2$ and $\varphi(T_2) \subseteq \mathbb{R}^2 \setminus \overline{G}$.

By relation $\varphi(w_1) \in G \subseteq \text{Int } D^2$ and Condition (2), the vertex w_1 has degree at least 2. Therefore it is adjacent to at least one edge of T except e that is an edge of T_1 . It means that a tree T_1 is non degenerated.

Since degrees of all other vertices of T_1 in T are the same as degrees in T_1 then $V_{\text{ter}}(T_1) \subseteq V_{\text{ter}}(T) \cup \{w_1\}$. As we know $\#V_{\text{ter}}(T_1) \geq 2$ whence there is $w \in V_{\text{ter}}(T_1) \cap V_{\text{ter}}(T)$.

By the construction $\varphi(w) \in G \subseteq \text{Int } D^2$.

On the other hand it follows from (1) and (2) that $\varphi(w) \in \varphi(V^*) \subseteq \partial D^2$.

We have the contradiction with the assumption that $U \setminus \varphi(T) \subseteq U_i$ for some i .

So, there are exactly two components $U_i \neq U_j$ of a compliment $\mathbb{R}^2 \setminus (\varphi(T) \cup \partial D^2)$ such that the point $x \in \varphi(e) \setminus \{\varphi(w_1), \varphi(w_2)\}$ which is contained in the image of an edge e of T is a boundary point of. Consequently, by Corollaries 1.1 and 1.2 there are exactly two paths such that they pass through an arbitrary edge of T and connect the adjacent vertices of V^* .

Let $\#V^* \geq 3$ and for any $e \in E(T)$ of T there are exactly two paths such that they pass through this edge and connect adjacent vertices of V^* .

We should prove that the tree T is $\mathcal{D}$ -planar.

At first we consider a relation C that is a full cyclic order on a set V^* . It generates a convenient relation ρ_C on V^* , see Definition 1.5. Let us examine a set of the directed paths

$$\mathcal{P} = \{P(v, v') \mid v' \rho_C v\}$$

in T .

By Definitions 1.1 and 1.5 two vertices $v, v' \in V^*$ are adjacent with respect to a cyclic order C iff either $v \rho_C v'$ or $v' \rho_C v$ is true. These correlations can not hold true simultaneously, since a pair of vertices v, v' would generate a ρ_C -cycle, see Definition 1.3, and this contradicts to Proposition 1.3 and Corollary 1.4 since $\#V^* \geq 3$.

It follows from the discussion above that for every edge e of T there are exactly two paths of the set $\mathcal{P}$ passing through e .

Let us consider a binary relation ρ on the set V_* which is defined by a correlation

$$(9) \quad v \rho v' \Leftrightarrow P(v, v') \in \mathcal{P}.$$

Evidently, relation ρ is *dual* to the relation ρ_C ($v_1 \rho v_2 \Leftrightarrow v_2 \rho_C v_1$). Therefore by Definition 1.2, ρ is the convenient relation on V^* . So, a minimal relation of equivalence $\hat{\rho}$ on V^* containing ρ coincides with a minimal relation of equivalence $\hat{\rho}_C$ on V^* containing ρ_C . Thus the elements of the set V^* generate a ρ -cycle, see Proposition 1.3 and Corollary 1.3.

Let e be an edge of the tree T . We should prove that those two directed paths of the set $\mathcal{P}$ that contain e pass through e in opposite directions.

Let us consider a binary relation μ_e on V^* that is defined as follows

$$v\mu_e v' \Leftrightarrow P(v, v') \in \mathcal{P} \text{ i } e \notin P(v, v').$$

It is easy to see that a diagram of the relation μ_e can be obtained from a diagram of ρ by removing two pairs of vertices of V^* corresponding to paths of $\mathcal{P}$ which pass through e . Let (v_1, v'_1) and (v_2, v'_2) be such pairs. Therefore the relation μ_e satisfies the conditions of Lemma 1.4.

By this Lemma a minimal relation of equivalence $\hat{\mu}_e$ containing μ_e has two classes of equivalence B_1, B_2 and $v_1 \in B_1, v_2 \in B_2$.

Let $w, w' \in V$ be the ends of e . Let us consider a subgraph T' of the tree T such that $V(T') = V(T)$ and $E(T') = E(T) \setminus \{e\}$. It is clear that the vertices w and w' belong to different connected components of a graph T' (if there exists a path P in T' such that it connects them then these vertices can be connected by two different paths P and $P' = \{e\}$ in the tree T). We denote these components by T_w and $T_{w'}$.

Suppose that for vertices $v, v' \in V$ there is an directed path $P(v, v')$ passing through e . Let it first passes through the vertex w and then through w' . Then paths $P(v, w)$ and $P(w', v')$ belong to T' , so $v \in T_w, v' \in T_{w'}$. In case when the path $P(v, v')$ first passes through w' and then through w we have $v' \in T_w$ and $v \in T_{w'}$.

It is easy to see that every class of equivalence of the relation $\hat{\mu}_e$ belongs to the unique connected component of the set T' . By the construction different classes of equivalence have to belong to the different components of T' .

So, we conclude that either $B_1 \subseteq T_w$ and $B_2 \subseteq T_{w'}$ or $B_1 \subseteq T_{w'}$ and $B_2 \subseteq T_w$. Suppose that first pair of inequalities holds true.

If the directed paths $P(v_1, v'_1)$ and $P(v_2, v'_2)$ pass through e in the same direction, then $P(v_1, w) \cup P(v_2, w) \subseteq T_w$ and $v_2 \in T_w$. By the construction $T_w \cap V^* = B_1$ thus $v_2 \in B_1$. But it is a contradiction to Lemma 1.4. So, the paths $P(v_1, v'_1)$ and $P(v_2, v'_2)$ pass through e in the opposite directions.

The case $B_1 \subseteq T_{w'}, B_2 \subseteq T_w$ is considered similarly.

Let us construct an embedding of T into oriented disk D^2 .

Let D^2 be an oriented disk (closed disk with a fixed orientation on the boundary), $I = [0, 1]$ an directed segment and $\psi : I \rightarrow D^2$ an embedding such that $\psi(I) \subseteq \partial D^2$. The direction of a segment is said to be *coordinated* with the orientation of disk if a direction of passing along the simple continuous curve $\psi(I)$ from the origin $\psi(0)$ to the end $\psi(1)$ coincides with given orientation of the boundary ∂D^2 .

Every directed path in T is topologically a closed segment thus for directed path $P(v, v')$ with the origin v and the end v' there exists an embedding $\Phi_{P(v, v')} : P(v, v') \rightarrow D^2$ such that $\Phi_{P(v, v')}(P(v, v')) \subseteq \partial D^2$ and a direction of $P(v, v')$ is coordinated with the orientation of D^2 .

We fix a disjoint union of closed oriented disks $\bigsqcup_{P \in \mathcal{P}} D_P$ and a set of the embeddings

$$(10) \quad \begin{aligned} \Phi_{P(v,v')} : P(v,v') &\rightarrow D_{P(v,v')}, \\ \Phi_{P(v,v')}(P(v,v')) &\subseteq \partial D_{P(v,v')}, \quad P(v,v') \in \mathcal{P}, \end{aligned}$$

such that the directions of paths $P(v,v') \in \mathcal{P}$ are coordinated with the orientations of corresponding disks.

Let us consider a space

$$\tilde{D} = T \sqcup \bigsqcup_{P \in \mathcal{P}} D_P.$$

All maps Φ_P , $P \in \mathcal{P}$ are injective therefore a family of sets

$$F_x = \begin{cases} \{x\} \cup \bigcup_{P \in \mathcal{P} : x \in P} \Phi_P(x), & x \in T, \\ \{x\}, & x \in \bigcup_{P \in \mathcal{P}} D_P \setminus \Phi_P(P). \end{cases}$$

generates a partition $\mathfrak{f}$ of the space $\tilde{D}$.

We consider a factor-space D of $\tilde{D}$ over partition $\mathfrak{f}$ and a projection map

$$\text{pr} : \tilde{D} \rightarrow D.$$

Let us prove that D is homeomorphic to a disk, the orientations of disks D_P , $P \in \mathcal{P}$ give some orientation on D and a map

$$\varphi = \text{pr}|_T : T \rightarrow D$$

conforms to the conditions of Definition 2.1.

At first we investigate some properties of the space D and the projection pr .

2.1. The mapping pr is closed.

Recall that a set is called *saturated* over partition $\mathfrak{f}$ if it consists of entire elements of that partition.

Topology of space D is a factor-topology (a set A is closed in D iff its full preimage $\text{pr}^{-1}(A)$ is closed in $\tilde{D}$). For proof of closure of a projection map pr it is sufficient to check that for any closed subset K of the space $\tilde{D}$ minimal saturated set $\tilde{K} = \text{pr}^{-1}(\text{pr}(K))$ containing K is also closed.

From the definition of partition $\mathfrak{f}$ it follows that

$$(11) \quad \begin{aligned} K &= (K \cap T) \sqcup \bigsqcup_{P \in \mathcal{P}} (K \cap D_P), \\ \tilde{K} &= (K \cap T) \sqcup \bigsqcup_{P \in \mathcal{P}} ((K \cap D_P) \cup \Phi_P(K \cap P)). \end{aligned}$$

Sets T , P , D_P , $P \in \mathcal{P}$ are compacts and all maps Φ_P are homeomorphisms onto their images. Thus all sets $K \cap T$, $K \cap D_P$, $\Phi_P(K \cap P)$, $P \in \mathcal{P}$, are compacts. The graph T is finite hence $\#\mathcal{P} < \infty$ and the union on the right of (11) is finite. The set $\tilde{K}$ is a compact, so it is closed.

We remark that we incidentally verified that the space $\tilde{D}$ is compact.

2.2. The space D is a compactum.

D is the compact space as a factor-space of compact space $\tilde{D}$. Compactum $\tilde{D}$ is the normal topological space and a factor-space of a normal space over closed partition is a normal space, see [4]. Thus D is a normal space, in particular, D is Hausdorff space. Therefore D is compactum.

2.3. Map $\varphi = \text{pr}|_T : T \rightarrow D$ is the embedding.

By definition, $F_x \cap T = \{x\}$ for every $x \in T$, hence φ is an injective map. The space T is compact and the space D is Hausdorff thus φ is homeomorphism onto its image, see [4].

2.4. For every $P \in \mathcal{P}$ a map $\text{pr}|_{D_P} : D_P \rightarrow D$ is an embedding.

By definition, for $x \in D_P$ we get

$$(12) \quad D_P \cap F_x = \begin{cases} \Phi_P(\Phi_P^{-1}(x)), & x \in \Phi_P(P), \\ \{x\}, & x \in D_P \setminus \Phi_P(P). \end{cases}$$

The map Φ_P is injective hence $\Phi_P(\Phi_P^{-1}(x)) = \{x\}$, $x \in \Phi_P(P)$. Finally, $F_x \cap D_P = \{x\}$ for every $x \in D_P$ and a continuous map $\text{pr}|_{D_P}$ is injective. Thus it is a homeomorphism of compact D_P onto its image.

2.5. For every $P \in \mathcal{P}$ a set $\text{pr}(D_P \setminus \Phi_P(P))$ is open in D and has no common points with a set $\text{pr}(\tilde{D} \setminus (D_P \setminus \Phi_P(P)))$.

Let $P \in \mathcal{P}$. The set D_P is open-closed in the space $\tilde{D}$, hence an open set $D_P \setminus \Phi_P(P)$ in D_P is also open in $\tilde{D}$. This set is saturated by definition. Therefore $D_P \setminus \Phi_P(P) = \text{pr}^{-1}(\text{pr}(D_P \setminus \Phi_P(P)))$ and a set $\text{pr}(D_P \setminus \Phi_P(P))$ is open in the factor-space D .

It follows from the discussion above that a set $\tilde{D} \setminus (D_P \setminus \Phi_P(P))$ is also saturated and it has no common points with $D_P \setminus \Phi_P(P)$. Thus

$$\text{pr}(D_P \setminus \Phi_P(P)) \cap \text{pr}(\tilde{D} \setminus (D_P \setminus \Phi_P(P))) = \emptyset.$$

2.6. Let $e \in E$ be any edge of the tree T , points w_1, w_2 be the ends of e and $P', P'' \in \mathcal{P}$ be paths in $\mathcal{P}$ that pass through e . We designate $e^0 = e \setminus \{w_1, w_2\}$,

$$\begin{aligned} D_{P'}^0 &= D_{P'} \setminus \partial D_{P'} \subseteq \bigcup_{p \in \mathcal{P}} D_p \setminus \Phi_p(P), \\ D_{P''}^0 &= D_{P''} \setminus \partial D_{P''} \subseteq \bigcup_{p \in \mathcal{P}} D_p \setminus \Phi_p(P), \\ \tilde{U} &= (D_{P'}^0 \cup \Phi_{P'}(e^0)) \sqcup (D_{P''}^0 \cup \Phi_{P''}(e^0)) \sqcup e^0, \\ U &= \text{pr}(\tilde{U}). \end{aligned}$$

U is the open neighborhood of a set $\text{pr}(e^0)$ in the space D , it is homeomorphic to open disk and is divided by a set $\text{pr}(e^0)$ onto two connected components $\text{pr}(D_{P'}^0)$ and $\text{pr}(D_{P''}^0)$.

To prove this we should remark that sets e^0 , $D_{P'}^0$ and $D_{P''}^0$ are open in $\tilde{D}$. By definition of partition $\mathfrak{f}$ for every $x \in e^0$ we get $F_x = \{x, \Phi_{P'}(x), \Phi_{P''}(x)\}$ since the set $\tilde{U}$ is saturated.

The set $\tilde{U}$ is open in $\tilde{D}$. Really, in the first place e^0 is an open subset of T , secondly, $\Phi_{P'}(e^0)$ is an open subset of closed subspace $\Phi_{P'}(P')$ of space $D_{P'}$, therefore, $\Phi_{P'}(P') \setminus \Phi_{P'}(e^0)$ is a closed subset $D_{P'}$. Let us remark that arcs $\Phi_{P'}(P')$ and $\partial D_{P'} \setminus \Phi_{P'}(P')$ have $\Phi_{P'}$ - images of endpoints of the path P' as common ends, thus $\overline{\partial D_{P'} \setminus \Phi_{P'}(P')} \cap \Phi_{P'}(e^0) = \emptyset$ and following conditions hold true

$$\begin{aligned} \partial D_{P'} \setminus \Phi_{P'}(e^0) &= (\partial D_{P'} \setminus \Phi_{P'}(P')) \cup (\Phi_{P'}(P') \setminus \Phi_{P'}(e^0)) = \\ &= \overline{(\partial D_{P'} \setminus \Phi_{P'}(P')) \cup (\Phi_{P'}(P') \setminus \Phi_{P'}(e^0))}. \end{aligned}$$

So, a set $\partial D_{P'} \setminus \Phi_{P'}(e^0)$ is closed in $D_{P'}$ and a set $D_{P'}^0 \cup \Phi_{P'}(e^0) = D_{P'} \setminus (\partial D_{P'} \setminus \Phi_{P'}(e^0))$ is open in $D_{P'}$. Similarly, a set $D_{P''}^0 \cup \Phi_{P''}(e^0)$ is open in $D_{P''}$. Sets T , $D_{P'}$ and $D_{P''}$ are open-closed in space $\tilde{D}$. Thus the set $\tilde{U}$ is open in $\tilde{D}$.

Finally, the set $U = \text{pr}(\tilde{U})$ is open in D . This set is a result of gluing

$$\begin{aligned} U &\cong (D_{P''}^0 \cup \Phi_{P''}(e^0)) \cup_{\alpha} (D_{P'}^0 \cup \Phi_{P'}(e^0)), \\ \alpha &= \Phi_{P''} \circ \Phi_{P'}^{-1} : \Phi_{P'}(e^0) \rightarrow \Phi_{P''}(e^0). \end{aligned}$$

A map α is a composition of homeomorphisms. Therefore U is homeomorphic to open disk and is divided by $\text{pr}(e^0)$ onto two connected components $\text{pr}(D_{P'}^0)$ and $\text{pr}(D_{P''}^0)$.

2.7. For any $P_1, \dots, P_n \in \mathcal{P}$ a boundary $\text{Fr } D_n$ of a set $D_n = \text{pr}(\bigcup_{i=1}^n D_{P_i})$ in the space D belongs to $\text{pr}(\bigcup_{i=1}^n P_i) = \text{pr}(T) \cap D_n$.

It follows from property 2.5 that $\text{Fr } \text{pr}(D_{P_i}) \subseteq \text{pr}(\Phi_{P_i}(P_i)) = \text{pr}(P_i)$ for any $i \in \{1, \dots, n\}$. Hence

$$\text{Fr } D_n \subseteq \bigcup_{i=1}^n \text{Fr } \text{pr}(D_{P_i}) \subseteq \bigcup_{i=1}^n \text{pr}(P_i) = \text{pr}\left(\bigcup_{i=1}^n P_i\right).$$

2.8. Let $P_1, \dots, P_n \in \mathcal{P}$, $\tilde{D}_n = \bigcup_{i=1}^n D_{P_i}$, $D_n = \text{pr}(\tilde{D}_n)$. Let e be an edge of T such that $\text{pr}(e^0) \cap D_n \neq \emptyset$, where e^0 is an edge e without ends.

A set $\text{pr}(e^0)$ belongs to $\text{Int } D_n$ iff at least one point $y \in \text{pr}(e^0)$ has a neighborhood in D_n which is homeomorphic to open disk. Otherwise, a set $\text{pr}(e^0)$ belongs to $\text{Fr } D_n$.

If $\text{pr}(e^0) \subseteq \text{Int } D_n$ then a set $\text{pr}(e^0)$ has a neighborhood in D_n which is homeomorphic to open disk and both paths $P', P'' \in \mathcal{P}$ passing through e belong to a set $\{P_1, \dots, P_n\}$.

If $\text{pr}(e^0) \subseteq \text{Fr } D_n$, then exactly one of them belongs to $\{P_1, \dots, P_n\}$.

Suppose that paths $P', P'' \in \mathcal{P}$ pass through the edge e . By the definition $\text{pr}(e^0) \subseteq D_n \cap \text{pr}(T) = \text{pr}(\bigcup_{i=1}^n P_i)$, so at least one of them belongs to $\{P_1, \dots, P_n\}$. We consider two possibilities.

We assume that $P' = P_k$, $P'' = P_s$, $k, s \in \{1, \dots, n\}$. Then a set

$$U = \text{pr}((D_{P'}^0 \cup \Phi_{P'}(e^0)) \cup (D_{P''}^0 \cup \Phi_{P''}(e^0)) \cup e^0) \subseteq D_n$$

is an open neighborhood of $\text{pr}(e^0)$ that is homeomorphic to an open disk, see 2.6.

Let $P' \in \{P_1, \dots, P_n\}$, $P'' \notin \{P_1, \dots, P_n\}$. In this case $U = U' \cup U'' \cup e^0$, $U' = \text{pr}(D_{P'}^0) \subseteq D_n$ but a set $U'' = \text{pr}(D_{P''}^0)$ has no common points with D_n , therefore $\text{pr}(e^0) \subseteq \text{Fr } D_n$.

Suppose that for some $y \in \text{pr}(e^0)$ in D_n there exists an neighborhood $W_y \in D_n$ homeomorphic to open two dimensional disk. By using theorem of Shenflies [11,13] we can find a small neighborhood $\hat{W}_y$ of y in D_n such that it is homeomorphic to an open disk and satisfies following conditions:

- A set $\overline{\hat{W}_y}$ in the space D_n is homeomorphic to a closed disk and is separated from compacts $\text{pr}(T \setminus e^0)$ and $D \setminus U$.
- $\overline{\hat{W}_y}$ intersects $\text{pr}(e^0)$ by a connected segment that is a cut of the disk $\overline{\hat{W}_y}$.

Then the set $\text{pr}(e^0)$ divides $\hat{W}_y$ onto two connected components $W_1 \cup W_2 = \hat{W}_y \setminus \text{pr}(e^0)$, $W_1 \cap W_2 = \emptyset$ such that $\overline{W_1} \cap \overline{W_2} \ni y$.

By the construction $\hat{W}_y \subseteq U \cap D_n$ and $W_1, W_2 \subseteq (U \cap D_n) \setminus \text{pr}(T)$. Let us remind that $P' \in \{P_1, \dots, P_n\}$, thus $\text{pr}(D_{P'}) = D' \subseteq D_n$. Similarly, $P'' \notin \{P_1, \dots, P_n\}$ hence $\text{pr}(D_{P''}^0) \cap D_n = \emptyset$, see property 2.5. Therefore $U \cap D_n = U' \cup \text{pr}(e^0)$, where $U' = \text{pr}(D_{P'}^0)$, see property 2.6, and $U \cap D_n \cap \text{pr}(T) = \text{pr}(e^0) \subseteq \partial D'$, where $D' = \text{pr}(D_{P'})$.

Thus $y \in \partial D'$ and the set $\hat{W}_y$ is the open neighborhood of y in closed disk D' and $\hat{W}_y \cap (D' \setminus \partial D') = W_1 \cup W_2$, $y \in \overline{W_1} \cap \overline{W_2}$. By Lemma 1.5 we can conclude that $W_1 = W_2$ but it contradicts to the assumption that $W_1 \cap W_2 = \emptyset$.

So, if $\{P', P''\} \not\subseteq \{P_1, \dots, P_n\}$, then there is no $y \in \text{pr}(e^0)$ that has an open neighborhood in D_n , which is homeomorphic to open disk.

2.9. Let $P_1, \dots, P_n \in \mathcal{P}$. Let us describe a structure of boundary $\text{Fr } D_n$ of $D_n = \text{pr}(\bigcup_{i=1}^n D_{P_i})$ in D .

Denote by $E_n \subseteq E$ a set of all edges of the tree T such that exactly one of two paths $P', P'' \in \mathcal{P}$ passing through $e \in E_n$ belongs to $\{P_1, \dots, P_n\}$. As we know, see Condition 2.8, $\text{pr}(E_n) \subseteq \text{Fr } D_n$ and if for some edge $e \in E$ we get $e \notin E_n$, then $\text{Fr } D_n \cap \text{pr}(e) \subseteq \{v', v''\}$, where $v', v'' \in V$ are ends of e .

Similarly, denote by $V_n \subseteq V$ a set of all vertices of T such that for a vertex $v \in V_n$ the following condition satisfies: $\text{pr}(v) \in D_n$ and all edges that are adjacent to v belong to $E \setminus E_n$. It is easy to show that the set V_n is discreet and $\text{pr}(E_n) \cap \text{pr}(V_n) = \emptyset$.

From the discussion above and Condition 2.7 it follows that

$$(13) \quad \text{pr}(E_n) \subseteq \text{Fr } (D_n) \subseteq (\text{pr}(E_n) \cup \text{pr}(V_n)).$$

2.10. Let $P_1, \dots, P_n \in \mathcal{P}$. A set $D_n = \text{pr}(\bigcup_{i=1}^n D_{P_i})$ is connected iff then $\bigcup_{i=1}^n P_i$ is a connected subgraph of the tree T .

Let $\bigcup_{i=1}^n P_i = T'$ is a connected subgraph of T . Then $D_n = \text{pr}(T') \cup \bigcup_{i=1}^n \text{pr}(D_{P_i})$, all sets $\text{pr}(T')$, $\text{pr}(D_{P_i})$, $i \in \{1, \dots, n\}$ are connected and $\text{pr}(T') \cap \text{pr}(D_{P_i}) \neq \emptyset$, $i \in \{1, \dots, n\}$. Hence the set D_n is connected.

Next, let $\bigcup_{i=1}^n D_{P_i} = T' \cup T''$, $T' \cap T'' = \emptyset$ and sets T' , T'' are nonempty and closed. Every set P_i , $i \in \{1, \dots, n\}$ is connected, therefore, either $P_i \in T'$ or $P_i \in T''$. Without loss of generality we can change indexing of the elements of $\{P_1, \dots, P_n\}$ in such way that for some $s \in \{1, \dots, n-1\}$ the following conditions are satisfied

$$T' = \bigcup_{i=1}^s P_i, \quad T'' = \bigcup_{i=s+1}^n P_i.$$

Every set

$$\tilde{D}' = T' \cup \bigcup_{i=1}^s D_{P_i}, \quad \tilde{D}'' = T'' \cup \bigcup_{i=s+1}^n D_{P_i},$$

is closed, whence sets $D' = \text{pr}(\tilde{D}')$ and $D'' = \text{pr}(\tilde{D}'')$ are closed, see Condition 2.1. By the construction $\tilde{D}' \cap \tilde{D}'' = \emptyset$. Let $y \in D' \cap D''$. A map pr is injective by definition on the set $\text{pr}^{-1}(D \setminus \text{pr}(T))$ and sets $\tilde{D}'$ and $\tilde{D}''$ do not intersect on $\text{pr}^{-1}(D \setminus \text{pr}(T))$, thus $y \in \text{pr}(T)$. Hence $y \in \text{pr}(T \cap \tilde{D}') \cap \text{pr}(T \cap \tilde{D}'') = \text{pr}(T') \cap \text{pr}(T'')$. But as we know, see Condition 2.3, the map $\varphi = \text{pr}|_T$ is bijective, therefore $\text{pr}(T') \cap \text{pr}(T'') = \text{pr}(T' \cap T'') = \emptyset$. We get a contradiction, thus $D' \cap D'' = \emptyset$.

Hence $D_n = D' \sqcup D''$ and sets D' , D'' are closed and nonempty. Therefore the set D_n is not connected.

Finally let us prove a $\mathcal{D}$ -planarity of the tree T .

Let for some n , $1 \leq n < \sharp \mathcal{P}$ directed paths $P_1 = P(v_1, v'_1), \dots, P_n = P(v_n, v'_n) \in \mathcal{P}$ are fixed and $\tilde{D}_n = \bigcup_{i=1}^n D_{P_i}$, $D_n = \text{pr}(\tilde{D}_n)$.

For every $i \in \{1, \dots, n\}$ we denote by $\tilde{\gamma}_i$ an directed arc of ∂D_{P_i} from point $\Phi_{P_i}(v'_i)$ to $\Phi_{P_i}(v_i)$ which has no other common points with an arc $\Phi_{P_i}(P_i)$.

Suppose that the objects under consideration comply with following conditions.

- (i) A space D_n is homeomorphic to a close two-dimensional disk.
- (ii) There exists at least one edge $e \in \bigcup_{i=1}^n P_i$ such that its image $\text{pr}(e)$ is contained in a boundary circle ∂D_n of D_n .
- (iii) A disk D_n is oriented in the following way: for every $i \in \{1, \dots, n\}$ and every edge $e \in P_i$ such that $\text{pr}(e)$ belongs to ∂D_n an orientation of e generated by the direction of $P_i = P(v_i, v'_i)$ maps by pr onto an orientation of D_n .
- (iv) For every $i \in \{1, \dots, n\}$ an arc $\gamma_i = \text{pr}(\tilde{\gamma}_i)$ connects a point $\text{pr}(v'_i)$ with a point $\text{pr}(v_i)$ and has no other common points with a set $\text{pr}(T)$ and orientation of this arc is consistent with the orientation of D_n .

We should remark that for $n = 1$ and any path $P = P_1 \in \mathcal{P}$ if we take an orientation on $D_1 = \text{pr}(D_P)$ induced from D_P by using pr , then Conditions (i)–(iv) always hold true. By the construction, Conditions (iii) and (iv) are true, (i) follows from Condition 2.4, (ii) follows from Condition 2.8.

We also remark that it follows from Condition 2.8 that an edge $e \in \bigcup_{i=1}^n P_i$ belongs to ∂D_n of D_n iff $e \in E_n$. Thus Condition (iii) is well-posed. As well all boundary points of D_n in the space D possibly except a finite number of isolated points from the set $\text{pr}(V_n)$ belong to ∂D_n .

Let an edge $e \in \bigcup_{i=1}^n P_i$ satisfies Condition (ii). Then $e \in E_n$ and there is the unique path $P_{n+1} = P(v_{n+1}, v'_{n+1}) \in \mathcal{P} \setminus \{P_1, \dots, P_n\}$ such that it passes through the edge e . Let $e \in P_l$, where $P_l \in \{P_1, \dots, P_n\}$ is the second path among two paths from the set $\mathcal{P}$ which passes through the edge e .

Let us consider a disk $D_{P_{n+1}}$ and its image $D' = \text{pr}(D_{P_{n+1}})$. By Condition 2.4 it is also the closed disk. Let $\Gamma = D_n \cap D'$. It is obvious that Γ is closed.

By Condition 2.5 a set $\text{pr}(D_{P_{n+1}} \setminus \Phi_{P_{n+1}}(P_{n+1}))$ is open in D and does not intersect D_n . It follows from Condition 2.4 that

$$\begin{aligned} \text{pr}(D_{P_{n+1}} \setminus \Phi_{P_{n+1}}(P_{n+1})) &= \text{pr}(D_{P_{n+1}}) \setminus \text{pr} \circ \Phi_{P_{n+1}}(P_{n+1}) = D' \setminus \text{pr}(P_{n+1}), \\ \text{pr}(\Phi_{P_{n+1}}(P_{n+1})) &= \text{pr}(P_{n+1}) \subseteq \overline{(D' \setminus \text{pr}(P_{n+1}))}. \end{aligned}$$

Therefore

$$\Gamma = \text{Fr } D_n \cap \text{Fr } D' \subseteq \text{pr}(P_{n+1}).$$

Let us apply Condition 2.9 to D_n and D' . By (13) the set Γ consists of images of edges which belong to the path P_{n+1} and possibly from a number of images of vertices of a tree T .

Let us check that the set Γ is connected.

If it is not the case it follows from what we said above that there are two vertices $w_1, w_2 \in V$, $w_1 \neq w_2$ of T such that they belong to the path P_{n+1} and a projection of a path $P(w_1, w_2) \subseteq P_{n+1}$ which connects them in T intersects Γ by a set $\{\text{pr}(w_1), \text{pr}(w_2)\}$. Then $\text{pr}(P(w_1, w_2)) \cap D_n = \{w_1, w_2\}$.

On the other hand, the set D_n is connected thus $T' = \bigcup_{i=1}^n P_i$ is a connected subgraph of T , see Condition 2.10. From Condition 2.7 it follows that $w_1, w_2 \in V(T')$, therefore there is a path $P'(w_1, w_2)$ connecting them in T' . This path has to connect w_1 with w_2 in T . But $\text{pr}(P'(w_1, w_2)) \subseteq D_n$ hence $P'(w_1, w_2) \neq P(w_1, w_2)$. So, vertices w_1 and w_2 of T can be connected in T by two different paths which is impossible in the tree T .

This contradiction proves that Γ is connected.

It follows from the connectedness of Γ and from the inclusion $\text{pr}(e) \subseteq \Gamma \cap \text{pr}(E_n)$ that $\Gamma \subseteq \text{pr}(E_n)$. Thus

$$\Gamma \subseteq \partial D_n \cap \partial D',$$

where $\partial D' = \text{pr}(\partial D_{P_{n+1}})$ is a boundary circle of the disk D' .

By the discussion above and from $\Gamma \subseteq \text{pr}(P_{n+1})$ it is easy to understand that

$$\Gamma = \text{pr}(P(v, v'))$$

for some $v, v' \in V \cap P_{n+1}$, $v \neq v'$.

It is obvious that $P(v, v')$ is homeomorphic to a closed segment. From the Conditions 2.3 and 2.4 it follows that it is embedded into a boundary circles ∂D_n and $\partial D'$ by means of maps

$$\begin{aligned}\psi_n &= \text{pr} \big|_{P(v, v')} : P(v, v') \rightarrow D_n, \\ \psi' &= \text{pr} \circ \Phi_{P_{n+1}} : P(v, v') \rightarrow D' .\end{aligned}$$

Therefore, a set

$$D_{n+1} = D_n \cup D' \cong D_n \cup_{\psi} D', \quad \psi = \psi_n \circ (\psi')^{-1},$$

is a result of a gluing of closed disks D_n and D' by a segment that is embedded into the boundary circles of these disks. Consequently the set D_{n+1} is homeomorphic to a closed disk.

Let us denote $\tilde{D}_{n+1} = \bigcup_{i=1}^{n+1} D_{P_i}$. It is clear that

$$D_{n+1} = \text{pr} \left(\bigcup_{i=1}^n D_{P_i} \right) \cup \text{pr}(D_{P_{n+1}}) = \text{pr} \left(\bigcup_{i=1}^{n+1} D_{P_i} \right) = \text{pr}(\tilde{D}_{n+1}).$$

Hence the space D_{n+1} constructed according to the set $\{P_1, \dots, P_{n+1}\}$ satisfies Condition (i).

Disks D_n and D' are oriented. The orientation of D' is generated by an orientation of $D_{P_{n+1}}$ by means of the map pr .

By Condition (iii) applied to D_n and D' we get two orientations on e . One of them is induced from an orientation of $P_l \supseteq e$ and is coordinated with orientation of D_n . Another is generated by direction of P_{n+1} and is consistent with an orientation of D' .

As we said above the directed paths $P_l, P_{n+1} \in \mathcal{P}$ containing an edge e have to pass through e in the opposite directions. Therefore the orientations induced on Γ from D_n and D' are opposite. Hence the orientations of D_n and D' are coordinated and generate an orientation of D_{n+1} . It complies with the following condition

- for any simple arc $\alpha : I \rightarrow \partial D_n \cap \partial D_{n+1}$ an orientation of α is consistent with orientation of D_{n+1} iff an orientation α is coordinated with orientation of D_n ;
- for any simple arc $\beta : I \rightarrow \partial D' \cap \partial D_{n+1}$ an orientation of β is consistent with an orientation of D_{n+1} iff it is coordinated with an orientation a disk D' .

Disks D_n and D' satisfy Conditions (iii) and (iv). So, according to what has being said D_{n+1} also satisfies Conditions (iii) and (iv).

Suppose that the set D_{n+1} does not satisfy Condition (ii). Then $E_{n+1} = \emptyset$, see Condition 2.9 and Remark (iii), and $\partial D_{n+1} \cap \text{pr}(T) \subseteq \text{pr}(V)$. Thus a set $\partial D_{n+1} \cap \text{pr}(T)$ is finite.

The following correlations are implicated from Condition 2.5

$$\partial D_{n+1} \setminus \text{pr}(T) \subseteq \text{pr}\left(\bigcup_{i=1}^{n+1} (D_{P_i} \setminus \Phi_{P_i}(P_i))\right) = \bigcup_{i=1}^{n+1} \text{pr}(D_{P_i} \setminus \Phi_{P_i}(P_i)).$$

From Condition 2.4 it follows that for every $i \in \{1, \dots, n+1\}$ a set $\text{pr}(D_{P_i} \setminus \partial D_{P_i}) \subseteq D_{n+1}$ is homeomorphic to an open disk. Hence

$$\bigcup_{i=1}^{n+1} \text{pr}(D_{P_i} \setminus \partial D_{P_i}) \subseteq D_{n+1} \setminus \partial D_{n+1}.$$

From this correlation it follows, see Condition (iv), that

$$\begin{aligned} \partial D_{n+1} \setminus \text{pr}(T) &\subseteq \left[\bigcup_{i=1}^{n+1} (\text{pr}(D_{P_i} \setminus \partial D_{P_i}) \cup \text{pr}(\partial D_{P_i} \setminus \Phi_{P_i}(P_i))) \right] \cap \partial D_{n+1} = \\ &= \bigcup_{i=1}^{n+1} \text{pr}(\partial D_{P_i} \setminus \Phi_{P_i}(P_i)) \subseteq \bigcup_{i=1}^{n+1} \text{pr}(\tilde{\gamma}_i) = \bigcup_{i=1}^{n+1} \gamma_i. \end{aligned}$$

A set $\bigcup_{i=1}^{n+1} \gamma_i$ is closed in D hence it is also closed in ∂D_{n+1} . Therefore, a set $\partial D_{n+1} \setminus \bigcup_{i=1}^{n+1} \gamma_i$ have to be an open subset of a space ∂D_{n+1} . But

$$\partial D_{n+1} \setminus \bigcup_{i=1}^{n+1} \gamma_i \subseteq \partial D_{n+1} \cap \text{pr}(T) \subseteq \text{pr}(V)$$

and this set is finite. Consequently,

$$\partial D_{n+1} = \bigcup_{i=1}^{n+1} \gamma_i.$$

From Condition (iv) it easily follows that open arcs $\gamma_i \setminus \{\text{pr}(v_i), \text{pr}(v'_i)\}$, $i \in \{1, \dots, n+1\}$ are pairwise disjoint. Therefore every point of a set $\partial D_{n+1} \cap \text{pr}(T) = \bigcup_{i=1}^{n+1} \{\text{pr}(v_i), \text{pr}(v'_i)\}$ is a common boundary point of exactly two arcs of the family $\{\gamma_i\}_{i=1}^{n+1}$.

It follows from the choice of an orientation of arcs γ_i , $i \in \{1, \dots, n+1\}$ that if for some $s, r \in \{1, \dots, n+1\}$ either $v_s = v_r$ or $v'_s = v'_r$ is true, then $s = r$. Thus for every $i \in \{1, \dots, n+1\}$ there is the unique $j(i) \in \{1, \dots, n+1\}$, such that $v_i = v'_j$ and if $r \neq s$ then $j(r) \neq j(s)$. We also remark that by the construction $n \geq 1$, thus $n+1 \geq 2$ and $j(i) \neq i$, $i \in \{1, \dots, n+1\}$.

Therefore, on the set $\{1, \dots, n+1\}$ there is a transposition σ without fix points such that $v_i = v'_{\sigma(i)}$, $i \in \{1, \dots, n+1\}$. Let $\sigma = c_1 \cdots c_k$ be a decomposition of σ into independent cycles. Let $c_1 = (i_1 \dots i_m)$. Then $v_{i_1} = v'_{i_2}, \dots, v_{i_{m-1}} = v'_{i_m}$, $v_{i_m} = v'_{i_1}$.

From the definition of the set $\mathcal{P}$ we get $v'_i \rho_C v_i$, $i \in \{1, \dots, n+1\}$, since $P_i = P(v_i, v'_i) \in \mathcal{P}$. So, it is true that

$$v_{i_1} \rho_C v_{i_2}, \dots, v_{i_{m-1}} \rho_C v_{i_m}, v_{i_m} \rho_C v_{i_1},$$

thus vertices of the set $M_1 = \{v_{i_1}, \dots, v_{i_m}\}$ generate a ρ_C -cycle, see Definition 1.3. From Corollary 1.4 it follows that the set M_1 is a class of equivalence of a minimal equivalence relation $\hat{\rho}_C$ which contains the relation ρ_C . By Proposition 1.3 and Corollary 1.4 the relation $\hat{\rho}_C$ has the unique class of equivalence V^* . Hence $M_1 = V^*$, $\sigma = c_1$, $n+1 = \sharp V^* = \sharp \mathcal{P}$ and $D_{n+1} = D$.

From what was said above it follows that for $n+1 < \sharp \mathcal{P}$ the disk D_{n+1} satisfies Condition (ii). Thus, for $n+1 < \sharp \mathcal{P}$ the disk D_{n+1} satisfies (i)–(iv), but for $n+1 = \sharp \mathcal{P}$ it complies with conditions (i) and (iv).

Finally, starting from any path $P = P_1 = P(v_1, v'_1) \in \mathcal{P}$, we can sort out elements of a set

$$\mathcal{P} = \{P_1 = P(v_1, v'_1), \dots, P_N = P(v_N, v'_N)\}$$

in a finite number of steps so that for every set

$$D_n = \text{pr} \left(\bigcup_{i=1}^n D_{P_i} \right), \quad n \in \{1, \dots, N-1\},$$

the conditions (i)–(iv) are true and for the set

$$D_N = \text{pr} \left(\bigcup_{i=1}^N D_{P_i} \right) = \text{pr} \left(\bigcup_{P \in \mathcal{P}} D_P \right) = \text{pr}(\tilde{D}) = D$$

conditions (i) i (iv) are also true.

Thus $D_N = D$ is closed oriented two-dimensional disk, $\varphi = \text{pr}|_T : T \rightarrow D$ is an embedding, see Condition 2.3.

For every edge $e \in E$ both paths of $\mathcal{P}$ passing through this edge belong to a set $\{P_1, \dots, P_N\}$, thus $E_N = \emptyset$ and $\partial D = \bigcup_{i=1}^N \gamma_i$ with open arcs $\gamma_i \setminus \{\text{pr}(v_i), \text{pr}(v'_i)\}$ are pairwise disjoint. It is clear that

$$\varphi(T) \cap \partial D = \bigcup_{i=1}^N \{\text{pr}(v_i), \text{pr}(v'_i)\} = \bigcup_{P(v, v') \in \mathcal{P}} \{\text{pr}(v), \text{pr}(v')\} = V^*.$$

An orientation of D generates some cyclic order O on the set $\text{pr}(V^*)$. A map $\varphi_0 = \varphi|_{V^*} : V^* \rightarrow \text{pr}(V^*)$ is bijective, therefore, a map φ_0^{-1} generates on the set V^* some cyclic order C' which is an isomorphic image of a cyclic order O ($C'(v_1, v_2, v_3) \Leftrightarrow O(\text{pr}(v_1), \text{pr}(v_2), \text{pr}(v_3))$).

We induce a convenient relation $\rho_{C'}$ on V^* , see Definition 1.5. From Condition (iv) it follows that for every $i \in \{1, \dots, N\}$ we have $v'_i \rho_{C'} v_i$. On the other hand, by definition of the set $\mathcal{P}$ it follows that $v' \rho_C v$ iff $P(v, v') \in \mathcal{P}$. But

$\mathcal{P} = \{P_1, \dots, P_N\}$, hence if $P(v, v') \in \mathcal{P}$, then $P(v, v') = P_i = P(v_i, v'_i)$ for some $i \in \{1, \dots, N\}$. Therefore the following conditions hold true

$$v' \rho_C v \Rightarrow v' \rho_{C'} v, \quad v, v' \in V^*,$$

and the relation $\rho_{C'}$ contains ρ_C .

With the help of convenient relations ρ_C and $\rho_{C'}$ we can induce on V^* the relations of cyclic orders C_{ρ_C} and $C_{\rho_{C'}}$, respectively, see Definition 1.6 and Proposition 1.4. From Definition 1.6 it easily follows that if $\rho_{C'}$ contains ρ_C then $C_{\rho_{C'}}$ contains C_{ρ_C} . In other words, an identical map Id_{V^*} is monomorphism of cyclic order C_{ρ_C} onto $C_{\rho_{C'}}$, see Definition 1.7. From Lemma 1.3 it follows that $C_{\rho_C} = C$ and $C_{\rho_{C'}} = C'$, hence the map Id_{V^*} is monomorphism of the cyclic order C onto C' . Lemma 1.2 implies that the map Id_{V^*} is an isomorphism of cyclic order C onto C' .

By the construction a map φ_0^{-1} is an isomorphism of cyclic order O onto C' thus φ_0 is an isomorphism of cyclic order $C = C'$ onto a cyclic order O which is induced onto $\varphi(V^*)$ from an oriented circle ∂D .

Finally, the map φ satisfies all conditions of Definition 2.1 and a tree T is $\mathcal{D}$ -planar.

□

REFERENCES

- [1] *Kuratovskiy K.* Topology. Vol. I. New edition, revised and augmented. Translated from the French by J. Jaworowski Academic Press, New York-London; Panstwowe Wydawnictwo Naukowe, Warsaw 1966 xx+560 pp.
- [2] *Kuratovskiy K.* Topology. Vol. II. New edition, revised and augmented. Translated from the French by J. Jaworowski Academic Press, New York-London; Panstwowe Wydawnictwo Naukowe, Warsaw 1966 xx+560 pp.
- [3] *Mel'nikov, O. V.; Remeslennikov, V. N.; Roman'kov, V. A.; Skorniyakov, L. A.; Shestakov, I. P.* General algebra. Vol. 1 (Russian) Mathematical Reference Library, "Nauka", Moscow, 1990. 592 pp.
- [4] *Fuks, D. B.; Rokhlin, V. A.* Beginner's course in topology. Geometric chapters. Translated from the Russian by A. Iacob. Universitext. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1984. xi+519 pp.
- [5] *Fuks, L.* Partially ordered algebraic systems (Russian) Translated from the English by I. V. Stel'leckiy. Edited by A. G. Kurosh Izdat. "Mir", Moscow 1965 342 pp.
- [6] *Yurchuk I.* Topological equivalence of functions of class $F(D^2)$ Zb. prac Inst. Math. NAS Ukraine, 2006. – V.3, №3. – 474-486 pp. (in Ukrainian)
- [7] *Y. Ikebe, M. Perles, A. Tamura and S. Tokunaga* The rooted tree embedding problem into points in the plane. Discrete Comput. Geom., 11(1994), 51-63 pp.
- [8] *Atsushi Kaneko, M. Kano* Straight line embeddings of rooted star forests in the plane. Discrete Applied Math., Vol.101 (2000), 167-175 pp.
- [9] *Norbert A'Campo* Planar trees, slalom curves and hyperbolic knots. Publications mathématique de l'I.H.É.S., No.88 (1998), 171-180 pp.
- [10] *Norbert A'Campo* Generic immersion of curves, knots, monodromy and gordian number. arXiv: math 9803081v2 [math. GT] 7 Apr 1999

- [11] *M. H. A. Newman* Elements of the topology of plane sets of points. Cambridge: Cambridge Univ. Press, 1964, 214 pp.
- [12] *Novak V.* Cyclically ordered sets. Czechoslovak Math.Journal.,1982. Vol. 32(107). P. 460-473.
- [13] *Zieschang H., Vogt E., Coldewey H.- D.* Surfaces and planar discontinuous groups. Springer-verlag, 1981.